

THE CRATO FOSSIL BEDS OF BRAZIL

Window into an Ancient World

Edited by

DAVID M. MARTILL

University of Portsmouth

GÜNTER BECHLY

Staatliches Museum für Naturkunde, Stuttgart

ROBERT F. LOVERIDGE

University of Portsmouth



CAMBRIDGE
UNIVERSITY PRESS

CAMBRIDGE UNIVERSITY PRESS
Cambridge, New York, Melbourne, Madrid, Cape Town, Singapore, São Paulo

Cambridge University Press
The Edinburgh Building, Cambridge CB2 8RU, UK

Published in the United States of America by Cambridge University Press, New York

www.cambridge.org
Information on this title: www.cambridge.org/9780521858670

© Cambridge University Press 2007

This publication is in copyright. Subject to statutory exception
and to the provisions of relevant collective licensing agreements,
no reproduction of any part may take place without
the written permission of Cambridge University Press.

First published 2007

Printed in the United Kingdom at the University Press, Cambridge

A catalogue record for this publication is available from the British Library

ISBN 978-0-521-85867-0 hardback

Cambridge University Press has no responsibility for the persistence or accuracy of URLs for external or third-party internet websites referred to in this publication, and does not guarantee that any content on such websites is, or will remain, accurate or appropriate.

Triangulifera but not with *Cretereisma*. Thus it appears that *Cretereisma* is the sister taxon of *Litophlebia*+*Triangulifera*. *Prottereisma* and *Misthodotes* were abundant in the Lower Permian while *Litophlebia* lived about 60 myr later. Hence *Cretereisma*, with a phylogenetic position between *Litophlebia* and the two Permian taxa, is the only known representative of an ephemeroid lineage which had retained many plesiomorphic features for about 110–160 myr.

11.4 Ephemeroptera: mayflies

Arnold H. Staniczek

Mayflies represent one of the basal branches of winged insects and have a world wide distribution. Stem group representatives of mayflies date back to the Carboniferous. More than 3,000 Recent species have been described.

The larvae of mayflies are obligatorily bound to freshwater habitats. The imagines of mayflies are short-lived insects with a life span that varies from a few hours to a few days. Mayflies are the only pterygote insects to retain more than one fully winged stage. The subimago is the alate penultimate stage and undergoes another moulting to become the imago. The subimaginal moulting is generally regarded as a vestigial adult moulting as it is present in the apterygote insect groups. The adult stages of mayflies are easily recognized by the presence of usually very long cerci, often accompanied by a median long terminal filament (paracercus). At rest the wings are folded vertically over the abdomen. The hind wings are considerably smaller than the forewings and can also be entirely lost in some taxa. The imaginal wings are generally translucent and glabrous, but the subimaginal wings usually have a greyish or milky tinge and are equipped with setae. The venation is characterized by the presence of multiple crossveins, a pronounced costal brace especially in the forewing, and the presence of vein MP. The wings generally retain a pronounced pleating and corrugation of the longitudinal veins. The adult stages of mayflies do not feed, and their mouth parts are atrophied. The antennae are very short and bristle-like, convergent on Odonata. Other features are connected with the aerial mating flight that the males perform above the water: the male eyes are generally larger than the female ones, in some taxa (Baetidae, Leptophlebiidae) they are extended to so-called turbinate eyes, which are also morphologically divergent. The male forelegs are elongated to grasp the females around the base of the forewing during copulation. The male claspers, modified abdominal leglets of abdominal segment VIII, additionally clutch the female abdomen during copulation. Males have paired penes, and females have paired gonopores, respectively. The female ovipositor is completely reduced, and only some taxa have developed a secondary egg guide. After mating the female mayfly simply drops its eggs into the water. The eggs are equipped with highly variable adhesive structures to prevent drifting.

The larvae of mayflies undergo numerous moults until they reach the final instar. Extant species usually bear seven pairs of tracheal gills, which insert laterally on the first seven abdominal segments. The gills are of variable shape but often specific for higher taxa. They can be either simple or may be composed of an upper and a lower branch. The gills can be shaped like a leaf (*Baetidae*), fringed (*Ephemeroidae*), finger-like in some *Leptophlebiidae*, miniaturized in the *Oligoneuriidae*, or may be located beneath a specialized gill that functions as a gill cover (*Caenidae*). In the *Heptageniidae* the upper gill portion is leaf-like, and the lower portion is represented by a gill tuft. The larvae are also characterized by the presence of the tail filaments. The chewing mouth parts of mayfly larvae are well developed. The secondary mandibular joint is not fixed as a ball-and-socket joint, and the mandible is able to perform gliding movements similar to the conditions in silverfish. The hypopharynx retains side lobes (*superlinguae*), and maxillary and labial palps are both three-segmented. Mayfly larvae mostly feed on detritus accumulated with their maxillae and mandibles. The filter-feeding *Setisura* even use specialized setae to filter their food out of the water currents with their forelegs. A few taxa (*Behningiidae*, *Ameletopsidae*) have carnivorous larvae. Mayflies are important elements of the freshwater food chain and the nutrient base for many carnivorous fish. The emergence of mayflies is often highly synchronized, and at times millions of specimens can be observed in dense swarms.

Systematics and phylogeny

The monophyly of mayflies is well established and supported by numerous apomorphic characters such as the Palmen body, the atrophy of mouth parts, and an air-filled gut with aerostatic function. Apart from the *Ephemeroptera*, there are only two other Recent basal lineages of pterygote insects, the *Odonata* and *Neoptera*. There is, however, no general agreement on the sister group of *Ephemeroptera*. A potential sister-group relationship between *Odonata* and *Ephemeroptera+Neoptera* is only substantiated by few putative synapomorphic characters, namely by the presence of a direct sperm transfer in *Ephemeroptera+Neoptera*. A monophyletic taxon *Paleoptera* (Martynov, 1925; *Ephemeroptera+Odonata*) has mainly been advocated by Kukalová-Peck (1985, 1987, 1991). Potential synapomorphies of the *Paleoptera* include an anal wing brace that is similarly developed in stem group representatives of both groups, a media with common stem, and the presence of intercalary veins. Other workers have proposed a sister-group relationship between *Ephemeroptera* and all other pterygote insects, the *Metapterygota* (Börner, 1909). The rigid secondary mandibular joint together with a different arrangement of mandibular musculature, the loss of the terminal filament and *superlinguae*, and the loss of the imaginal moults are regarded as apomorphic characters of the

Metapterygota (see Kristensen, 1995 and Staniczek, 2000 for a detailed discussion). Molecular data are inconclusive (Ogden and Whiting, 2003), and a final decision cannot be made at present.

The phylogenetic systematics of Ephemeroptera is far from being fully resolved and remains controversial with regard to the composition and branching sequence of many higher taxa. There are about 37 monophyletic taxa recognized that are categorized as families. McCafferty and Edmunds (1979) divided the Ephemeroptera into the paraphyletic Schistonota and the Pannota, which are defined by medially fused larval wing pads. Later Kluge (in an oral conference contribution in 1992; and 1998) and then McCafferty (1997) excluded the Batiscidae+Prosopistomatidae from the Pannota and assumed the Prosopistomatoidea (*Carapacea sensu* McCafferty, 1997; *Posteritorna sensu* Kluge, 1998) to be the sister group of all other mayflies. Other similarities in both systematics are a proposed taxon of several rather plesiomorphic families, the Tridentiseta (Kluge, 2004) or Pisciforma (McCafferty, 1998), that consist of the Siphlonuroidea (Kluge *et al.*, 1995) and Baetoidea (Staniczek, 1997). There is also congruence in uniting all filter-feeding taxa with the Heptageniidae to a taxon Setisura (McCafferty, 1997) or Branchitergaliae (Kluge, 2004). The remaining groups, united into a taxon Furcatergaliae by Kluge (2004), are also considered to be monophyletic. Within these higher taxa, however, the arrangement of the different families differs considerably. Moreover, recent molecular systematics by Ogden and Whiting (2005) failed to support most of these groupings.

Evolution

One of the most important steps in insect evolution was the acquisition of wings. The development of insect wings from thoracic side lobes, the paranota, has been the prevailing hypothesis for a long time (Snodgrass, 1935), until Kukalová-Peck (1978, 1983) publicized the development of insect wings from modified thoracic gills. Both hypotheses claim to gain support from fossil evidence. The Recent ephemeropteran lineage is only one of several branches of early winged insects. The †Paleodictyopterida comprise several distinctive groups, all of which became extinct by the end of the Permian. At present there are some doubts whether the †Paleodictyopterida are monophyletic, as often assumed, because of the characteristic elongated, sucking mouth parts they share. In several groups of these Palaeozoic insects, huge prothoracic side lobes were developed as winglets. These prothoracic paranotal lobes had a similar venation to the meso- and metathoracic wings. The presence of prothoracic winglets in many of these Palaeozoic insects is generally taken as support for the above-mentioned paranotal hypothesis. However, paranota are rigid outgrowths that do not have any articulation with the prothorax. The

gill hypothesis assumes that insect wings are serially homologous with the gills of mayfly larvae (Kukalová-Peck, 1991). In particular, gills and wing pads of Permian larvae of †*Prottereismatidae* look similar to each other, and the wing pads, unlike in modern mayfly larvae, seem to articulate with the thorax. Consequently, an aquatic origin of pterygote insects has been postulated by the advocates of the gill hypothesis. On the other hand, all primarily apterous insect groups are terrestrial, and the basal taxa of ptergotine insects with aquatic larvae (Ephemeroptera, Odonata, Plecoptera) have developed very different modes of respiration. This makes it unlikely that the pterygotes evolved in an aquatic environment (Pritchard *et al.*, 1993), although there are no terrestrial larvae of stem group representatives of mayflies known. Recently the gill hypothesis has been modified and the use of wings as a skimming device has been proposed as the trigger of wing development (Marden and Thomas, 2005). Given the fact that skimming in Recent mayflies and stoneflies only occurs in a few derived taxa, this scenario is not very likely.

However, notable differences from early stem group representatives of mayflies to modern Ephemeroptera can be observed. While a costal brace is present in most fossil and modern taxa, its basal course is slightly different in early groups such as the †*Prottereismatidae*, where the costal brace is still well separated from the costal vein. Fore- and hind wings in the first fossil taxa are of the same size. Heteronomous wings first appear in the Jurassic, and all modern mayflies have significantly reduced hind wings. Other differences concern the development of a wing tornus in modern mayflies, while the hind margin of the wing in Palaeozoic stem group representatives has no sharp angle (Kluge, 2004). Generally the size of mayflies diminished during their evolution. Adults of Palaeozoic stem group representatives like the †*Prottereismatidae* are also assumed to have functional mouth parts (Grimaldi and Engel, 2005). The articulated wing pads of the associated larvae grew out in a laterocephalad direction. In modern mayflies the wing pads are fused with the mesonotum and grow parallel to the longitudinal body axis. The amount of gill numbers has become reduced from nine pairs to seven.

Fossil record

Winged insects were already abundant in the Late Cretaceous, but the discovery of the Early Devonian *Rhyniognatha hirsti*, a species that may belong within the lineage of Metapterygota, suggests that the age of the Metapterygota and of its putative sister group, the Ephemeroptera, dates back to that time (Engel and Grimaldi, 2004). The first fossils that have been assigned to the ephemeropteran lineage date back to the Late Carboniferous. While the assignment of two enigmatic fossils, *Triblosoba pulchella* from the Late Carboniferous of Commentry and *Lithoneura lameerei* from

the Late Carboniferous of Mazon Creek, to the stem group of mayflies is controversial (Willmann, 1999), most researchers agree on the assignment of *Bojophlebia prokopi*, from the Upper Carboniferous of Bohemia, to the stem group of mayflies (Kukalová-Peck, 1985). With a wing span of 45 cm it is also the largest mayfly precursor ever found.

There are many fossil stem group representatives from the Permian fauna preserved, among them the well-known †*Prottereismatidae* from the Early Permian of Kansas, Oklahoma, Moravia and the Urals (Tillyard, 1932; Carpenter, 1979; Hubbard and Kukalová-Peck, 1980). Another well-known taxon is the †*Misthodotidae* that together with the †*Prottereismatidae* disappeared by the end of the Permian (Tshernova, 1965). The stem group representatives were often placed in a separate suborder, ‘†*Permoplectoptera*’, which is, however, regarded as paraphyletic with respect to the crown group. Most of these early taxa went extinct by the end of the Permian, but others could be recorded from the Triassic (Sinitshenkova *et al.*, 2005). The youngest stem group representatives of Ephemeroptera have been found within the Crato Formation and are described in detail in the preceding Section 11.3, by Willmann. A further, undescribed, stem group mayfly from this locality is featured in Figures 11.90i and j. For recent reviews on all stem group representatives of mayflies, see Kluge (2004) and Willmann (this volume, Section 11.3).

The first species that are positioned within the crown group of Ephemeroptera appear in the Lower Jurassic (Tshernova, 1967): *Stackelbergisca sibirica* turned out to be closely related to the Recent *Siphluriscus chinensis* (Siphluriscidae; Zhou and Peters, 2003). Species that are attributed to the Leptophlebiidae can be traced back to the Jurassic (Hubbard and Savage, 1981). Numerous other Jurassic genera have been placed in the extinct taxa †*Epeoromimidae*, †*Aenigmephemeridae*, †*Mesephemeridae* and †*Hexagenitidae* (McCafferty, 1990). The affinities to Recent mayfly groups remain mostly uncertain (Kluge, 2004).

The Cretaceous mayfly fauna apart from the Crato Formation has been documented by fossil records from Transbaykalia (Tshernova and Sinitshenkova, 1974; Sinitshenkova, 1986), China (Lin and Huang, 2001), England (Sinitshenkova and Coram, 2002), Australia (Jell and Duncan, 1986) and Algeria (Sinitshenkova, 1975).

The oldest mayflies in amber have been recorded from the mid-Cretaceous of Myanmar (Sinitshenkova, 2000), Lebanon (McCafferty, 1997) and New Jersey (Peters and Peters, 2000; Sinitshenkova, 2000). Several mayfly fossils from Baltic and Dominican amber have been recorded from the Eocene and Miocene that mostly reflect the modern fauna (Staniczek and Bechly, 2002; Staniczek, 2003; Godunko, 2004; Godunko and Neumann, 2006). New families of mayflies are defined only rarely, which, however, can all be attributed to Recent monophyletic groups (Kluge

et al., 2006). Kluge (1986) even reports a putative Recent species from Baltic amber.

Palaeobiology and palaeoecology

Mayflies are obligatorily bound to fresh water. Extant species of mayfly are generally halophobic and only a few species are reported to tolerate elevated salt concentrations as they are present in brackish water (Chadwick *et al.*, 2002). *Protoligoneuria limai* (Figures 11.4a and 11.9c; Plates 7i and 8a–c) is reported to be the most common fossil insect species of the Crato Formation (Martins-Neto, 1996a; D. Martill, personal communication), although mayflies only make up about 7% of the described Crato species (Bechly, 1998c). The mass occurrence of thousands of fossil larval mayflies in the Crato limestones clearly points to the presence of streams in the immediate vicinity of the deposit. McCafferty (1990) and Tshernova (1970) hypothesized quiet waters as a habitat for *P. limai* because of its larval swimming adaptations such as the minnow-like body shape, siphlonuroid caudal appendages and its enlarged seventh gill. The larvae of the burrowing Ephemeroidea (Figures 11.4c and g, 11.5a and b and 11.6h and i; Plate 8f) found in Crato also point to the presence of streams or at least stream sections with a rather low flow rate. However, a rheophilic habitat cannot entirely be ruled out. Similar to the Odonata fauna with a high percentage of Gomphidae found here, the oligoneuriid species found were probably more adapted to a lotic environment and generally depended on a high amount of oxygen. Most probably the Crato environment provided a diverse habitat suitable for both lotic and lentic species. The few adult records compared to the enormous amount of larval specimens found may be explained by two factors: mayfly larvae are sensitive to downstream drift, and this is counterbalanced by an upstream compensation flight of the female adults. Both effects may lead to a spatial separation of larvae and adults with the effect of an accumulation of larvae which have been swept into the area of deposition.

Crato fossils

The first fossil mayfly from this locality was reported by Costa-Lima (1950) and described by Demoulin (1955). According to Martins-Neto (2005b) there have been 15 species of fossil Crato mayflies described, not including the two species of stem group mayflies described by Willmann in Section 11.3 (a further undescribed specimen is featured on Figures 11.90i and j). Important contributions to Crato mayflies were made by McCafferty (1990) and Martins-Neto (1996a). Shorter contributions were provided by Brito (1987), Martins-Neto and Caldas (1990), Polegato and Zamboni (2001), Zamboni (2001) and Bechly *et al.* (2001a: figures 28, 30, 36 and 37).




Fig. 11.4. Crato Formation Ephemeroptera: (a) Hexagenitidae, *Protoligoneuria limai*, composite reconstruction of larva (after McCafferty, 1990: figure 1); (b) Oligoneuriidae, *Colocrus indivicum*, reconstruction of forewing, adult paratype AMNH 43499 (after McCafferty, 1990: figure 19); (c) Ephemeroidae (Potamanthidae?), *Olindinella gracilis*, larval holotype AMA-I-028 (after Martins-Neto and Caldas 1990: figure 2); (d) Euthyplociidae, *Pristiplocia rupestris*, forewing, adult holotype AMNH 44308 (after McCafferty, 1990: figure 23); (e) Ephemeroidae, *Australiephemera revelata*, forewing and hind wing fragment, composite from adult holotype AMNH 44300 and adult paratype AMNH 44310 (after McCafferty, 1990: figures 20 and 21); (f) Ephemeridae, *Microephemera neotropica*, forewing, adult holotype AMNH 43303 (after McCafferty, 1990: figure 22); (g) Ephemeroidae (Ephemeridae?), *Cratonympha microcelata*, larval holotype GP/1T-1677 (after Martins-Neto and Caldas 1990: figure 1A).




Fig. 11.5. Crato Formation Ephemeroptera: (a) Ephemeroidea indet., larva, SMNS 66630; (b) Ephemeroidea indet., larva, coll. MSF; (c) Ephemeroidea indet., adult, coll. MSF; (d) Leptophlebiidae (?) sp. 1 (*sensu* McCafferty 1990), subimago?, SMNS 66639; (e) Familia incertae sedis, gen. et sp. nov., larva Z2 coll. MSF; (f) Familia incertae sedis, gen. et sp. nov., larva, SMNS 66622. Scale bars, 5 mm.

†Hexagenitidae Lameere, 1917

Apart from the Crato fossils, there are five more genera that have been placed in the extinct †Hexagenitidae: *Hexagenites* Scudder 1880, *Ephemeropsis* Eichwald 1864, *Hexameropsis* Tshernova and Sinitshenkova 1974, *Mongologenites* Sinitshenkova 1986 and *Caenoephemera* Lin and Huang 2001. The first mayfly ever from Crato was mentioned by Costa-Lima (1950), who assigned three larval specimens to the Baetidae (equivalent to today's 'Siphlonuroidea'); however, no formal description was given. Based on these specimens, Demoulin (1955) formally named the species as *Protoligoneuria limai*, and placed it in the Oligoneuriidae. Brito (1987) described a new genus and species *Palaeobaetodes limai* but McCafferty (1990) synonymized Brito's species with *Protoligoneuria limai*. He placed the species in the †Hexagenitidae, based on the characteristic, significantly enlarged and out-spread seventh gill that is also present in other hexagenitid larvae. This enlarged seventh gill is regarded as an autapomorphic larval character of the †Hexagenitidae




Fig. 11.6. Crato Formation Ephemeroptera: (a) Oligoneuriidae, *Colocrus magnum* sp. nov., putative adult, SMNS 66623; scale bar, 10 mm; (b) Oligoneuriidae, *Colocrus magnum* sp. nov., left wings of adult, SMNS 66623; (c) Euthyplociidae, *Pristiplocia rupestris*, adult, SMNS 66541; scale bar, 5 mm; (d) Euthyplociidae, *Pristiplocia rupestris*, right wings of adult, SMNS 66541; (e) Euthyplociidae, *Pristiplocia* sp., larva, SMNS 66539; scale bar, 5 mm; (f) Euthyplociidae, *Pristiplocia* sp., larva, SMNS 66621; scale bar, 3 mm; (g) Baetiscidae, *Protobaetisca bechlyi* gen. et sp. nov., larval holotype SMNS 66620, scale bar, 3 mm; (h) Ephemeroidea indet., larva, MURJ without number; scale bar, 3 mm; (i) Ephemeroidea indet., larva, SMNS 66634; scale bar, 3 mm.

(Kluge, 2004), but according to McCafferty (1990) it may only be a synapomorphic character between some taxa within the †Hexagenitidae. Martins-Neto (1996a) reinstated *Palaeobaetodes* and described a second species, *Palaeobaetodes britoi*, from larvae, and also described a third genus *Cratogenites*, including *C. corradiniae*, from larvae.

The known imagines of the †Hexagenitidae all have a characteristic wing pattern with a branched CuA in the forewing and several successive triads present between CuA₁ and CuA₂. The first adult specimen of †Hexagenitidae from Crato was described as yet another new genus and species, namely *Cratogenitoides delclosi*, by Martins-Neto (1996a).

Protoligoneuria limai Demoulin, 1955

Baetidae (Siphlonurinae), Costa-Lima, 1950: 419

Protoligoneuria limai Demoulin, 1955: 271. (Figures 11.4a and 11.9c; Plates 7i and 8a–c)

Paleobaetodes costalima Brito, 1987. Restored status as Syn. (Figure 11.7a)

Siphgondwanus occidentalis McCafferty, 1990 Syn. nov. (Figure 11.7h)

Palaeobaetodes britoi Martins-Neto, 1996a Syn. nov. (Figures 11.7c–d)

Cratogenites corradiniae Martins-Neto, 1996a Syn. nov. (Figures 11.7e–f)

Cratogenitoides delclosi Martins-Neto, 1996a Syn. nov. (Figure 11.8b)

Material: larval holotype specimen CD 6616, Vulcano collection (according to Martins-Neto (2005a) it is the only specimen that is left from the original three syntypes, and thus it is here designated as lectotype); more than 2,000 specimens deposited in the collection Martins-Neto, São Paulo; 77+4 larval specimens deposited at AMNH, and specimen AMNH 43404 (holotype of *Siphgondwanus occidentalis*); 236 larval specimens deposited at SMF, including the two specimens with preliminary nos Q4 and Q9 (Plates 7i and 8a); three very well-preserved specimens SMNS 66536, SMNS 66537 (Figure 11.9c) and SMNS 66672; and numerous specimens in other collections. Two putative alate specimens are no. RGZN-T002 (holotype of *Cratogenitoides delclosi*) in the Martins-Neto collection at the University of São Paulo, and specimen SMNS 66635 (Figures 11.9a and b).

Diagnosis: putative alate stages (Figures 11.9a and b) with compact anteritornous wing, relation of wing length to width of about 2:1. Length of forewing about 18–20 mm. C, Sc and R₁ not branched; Sc not shortened. Costal field and entire wing with numerous crossveins. RS basally branched at about one-quarter of wing length. MA branched in apical half of wing at about three-quarters of wing length. MP basally branched, with very short common stem, CuA basally branched at about one-third of distance between wing base and tornus. CuA field with four successive ‘hexagenitid’ triads. CuP not branched. A₁ with common short stem and basally branched. Hind wing significantly reduced, without costal process, with RS, MA and MP apically branched. No conspicuous gemination of longitudinal veins.




Fig. 11.7. Crato Formation Ephemeroptera (invalid taxa): (a, b) *Palaeobaetodes costalimai* Brito, 1987 (= *Protoligoneuria limai*); (a) mature nymph; scale bar, 5 mm, (b) alar; (c, d) *Palaeobaetodes britoi* Martins-Neto, 1996a (= *Protoligoneuria limai*); (c) head and thorax, RGMN-T006; (d) alar; (e, f) *Cratogenites corradiiniae* Martins-Neto, 1996a (= *Protoligoneuria limai*), holotype, RGMN-T001; (e) mature larva; scale bar, 1 mm; (f) head and thorax; (g) *Cratoligoneuriella leonardii* Martins-Neto, 1996a (= *Colocrus indivicum*), holotype nymph, RGMN-T005; (h) *Siphgondwanus occidentalis* McCafferty, 1990 (= *Protoligoneuria limai*), larva in dorsal view, legs omitted; scale bar, 1 mm. (a–h) After Martins-Neto (1996a); (i) after McCafferty (1990).




Fig. 11.8. Crato Formation Ephemeroptera: (a) *Ephemeroptera incertae sedis*, *Costalimella nordestina* Martins-Neto, 1996a, holotype, RGMN-T004; (b) *Cratogenites delclosi* Martins-Neto, 1996a (synonym of *Protoligoneuria limai*), holotype, RGMN T002. Scale bars, 5 mm. After Martins-Neto (1996a).

Larvae (Figures 11.4a and 11.9c; Plates 7i and 8a–c): body length up to 16 mm, excluding cerci. Siphlonuroid minnow-like appearance (Figures 11.4a and 11.9c), cerci at their inner sides and terminal filum with fringes of swimming hairs, cerci slightly longer than terminal filament, abdomen with seven pairs of tracheal gills, gill VII distinctively longer than other gills, all gills consisting of a leaf-like upper part and a lower gill tuft (Plates 7i and 8a) that previously had been overlooked. Anterior margin of gills I–VII strengthened, gill VII (Plate 8b) also with a second longitudinal rib in the posterior half of the gill, well apart from its posterior strengthened margin (often a small but distinctly protruding apical lobe is visible at the posterior end of this gill plate). Wing pads medially partially fused as in Recent species of *Setisura* (Plate 8c).

Comments: Brito (1987) described with *Palaeobaetodes costalima* (Figures 11.7a and b) a new genus and species of †Hexagenitidae from Crato. McCafferty (1990) synonymized *Palaeobaetodes* with *Protoligoneuria*, but Martins-Neto (1996a) reinstated the latter genus and described a second species, *Palaeobaetodes britoi* (Figures 11.7c and d). However, these authors mention only few differences to *Protoligoneuria*: according to Martins-Neto (1996a), *Paleobaetodes* differs from *Protoligoneuria* mainly in larger eyes, which occupy 50% of the head in *Palaeobaetodes* instead of 30% in *Protoligoneuria*. Obviously the author did not consider the sexual dimorphism of the eyes in many mayfly species. It is very common that male larvae have larger eyes than the females, and this has to be assumed in this case as well. A different ratio of gill lengths VII/VI was claimed to be the significant difference between the two described species of *Paleobaetodes* (Martins-Neto, 1996a), but I could not confirm a significant grouping in the material available (several hundred larvae). Consequently, both species of *Paleobaetodes* are here regarded as junior synonyms of *Protoligoneuria limai*.




Fig. 11.9. Crato Formation Ephemeroptera: (a, b) Hexagenitidae, *Protoligoneuria limai* Demoulin, 1955, wings of adult, SMNS 66635; (a), left wings; (b) right wings; (c) *Protoligoneuria limai*, larva, SMNS 66537; (d) *Cratohexagenites longicercus* gen. et sp. nov., larva, holotype no. 447 MURJ; (e) *Cratohexagenites longicercus* gen. et sp. nov., putative adult, no. O46 coll. MSF; (f) *Cratohexagenites minor* gen. et sp. nov., larva, holotype MB.I.2026; (g) Oligoneuriidae, *Colocrus indivicum*, larva, SMNS 66538; (h) Oligoneuriidae, *Colocrus indivicum*, larva, MB.I.2025; (i) Oligoneuriidae, *Colocrus magnum* sp. nov., larva, holotype, SMNS 66624.

Martins-Neto (1996a) also described *Cratogenites corradiniae* (Figures 11.7e and f). Apart from larger eyes the only differences to *P. limai* mentioned are longer antennae (about twice the head width) in *C. corradiniae*. However, while the line drawing of the holotype (Martins-Neto, 1996a: figure 2A) shows an elongated left antenna, the corresponding photograph (Martins-Neto, 1996a: Plate 1, figure 1) does not support this view. Hence I also synonymize *Cratogenites corradiniae* with *Protoligoneuria*, until there is sound proof that it can indeed be distinguished as a separate species from *P. limai*.

Martins-Neto also noted differences in the larval head shape of the specimens he investigated and tried to link these differences with different species. However, it seems more likely to me that the observed differences are due to different positions of the head. I also noted slight differences in the body shape, but this could be caused by a different state of nutrition or by a different phase of the moulting period.

The body length of *Cratogenitoides delclosi* (Figure 11.8b), the only hexagenitid species described in the alate state, closely matches the length of the largest larvae of *P. limai* found. So *C. delclosi* is regarded as a junior synonym of *P. limai* herein. A very well-preserved adult specimen is SMNS 66635 (Figures 11.9a and b).

None of the previous authors observed ventral gill tufts in the larvae they investigated. This may be due to the fact that the gill tufts are inconspicuous and only visible in very well-preserved specimens. However, once I had encountered well-preserved specimens with clearly visible gill tufts, they could be observed quite frequently in many specimens (Plates 7i and 8a). In some specimens the fibrillous endings of the tufts just slightly protrude the rear margin of the upper gill portion and give the leaf-like gills the appearance of having a fringed rear margin. McCafferty (1990) described a single siphlonurid larva as *Siphgondwanus occidentalis* (Figure 11.7h) having exactly the same gill shape as mentioned above. Furthermore, the seventh pair of gills is not well preserved in this specimen. It is very probable that this in fact is nothing but a badly preserved specimen of *P. limai*. Hence *Siphgondwanus occidentalis* is also synonymized with *Protoligoneuria limai*.

Demoulin (1955) assigned *P. limai* to the Oligoneuriidae, but McCafferty (1990) demonstrated that there are larval and imaginal apomorphies present that clearly point to a position of *P. limai* within the †Hexagenitidae.

The phylogenetic relationships of the †Hexagenitidae have been enigmatic for a long time. Tshernova and Sinitshenkova (1974) maintain affinities between the †Hexagenitidae and Siphlonuridae, but the characters they have in common are generally thought to be plesiomorphic. Demoulin (1971) emphasized similarities between the basal Recent oligoneuriid genus *Chromarcys* and the †Hexagenitidae, assuming a closer phylogenetic relationship between these taxa. However, *Chromarcys* is generally regarded as the sister group to all other Recent Oligoneuriidae (McCafferty, 1991), and the †Hexagenitidae are clearly lacking the autapomorphies

of the Oligoneuriidae, so the †Hexagenitidae cannot be an ingroup taxon of the Oligoneuriidae. If the gill tufts now found in *P. limai* should be confirmed as ground-plan characters of the †Hexagenitidae, a position of the †Hexagenitidae within the Setisura (McCafferty, 1991) = Branchitergaliae (Kluge, 2004), or a sister-group relationship between the two taxa, would be likely. However, a detailed structural analysis of the gill tufts is yet to be undertaken. A second character also supports this hypothesis: in some very well-preserved specimens a partial medial fusion of the wing pads is visible (Plate 8c). The same character state is also present in the filter-feeding taxa of the Branchitergaliae.

Cratohexagenites gen. nov.

Type species: *C. longicercus* sp. nov., by present designation.

Derivation of name: named after the type locality and the genus *Hexagenites*.

Diagnosis: Body of broader drop-like shape; seventh gill distinctly larger in proportion to the other gills than in *Protoligoneuria*, and of trapezoidal shape with angular instead of rounded hind margin.

Cratohexagenites longicercus sp. nov.

Material: larval holotype (Figure 11.9d) no. 447 (old no. G76) in coll. MURJ; larval paratype MB.I.2026 at MNB; putative adult specimen no. MSF O46 (Figure 11.9e), deduced from the corresponding large size of this imaginal hexagenitid.

Type locality: Chapada do Araripe, vicinity of Nova Olinda, southern Ceará, north-east Brazil.

Type horizon: Lower Cretaceous, Upper Aptian, Nova Olinda Member of the Crato Formation.

Derivation of name: named after its elongated caudal filaments.

Diagnosis and description (Figure 11.9d): hexagenitid larva with enlarged seventh gill of trapezoid shape, differs from *P. limai* by body length of about 27 mm, length of cerci 28 mm, length of terminal filament 24 mm. General body shape is drop-like. Length of gills I–VI is about 5 mm, length of gill VII is 11 mm. Otherwise quite similar to *P. limai*.

Comment: adult specimen no. MSF O46 has the characteristic CuA branch and apical MA branch of †Hexagenitidae (CuA field not preserved). Its body length of 23.5 mm corresponds to the large size of the holotypical larva. Like *Protoligoneuria* and Recent Setisura this new taxon also has the larval wing pads medially partially fused (Figure 11.9d).

Cratohexagenites minor sp. nov.

Material: larval holotype (Figure 11.9f) no. MB.I.2026 at MNB.

Type locality: Chapada do Araripe, vicinity of Nova Olinda, southern Ceará, north-east Brazil.

Type horizon: Lower Cretaceous, Upper Aptian, Nova Olinda Member of the Crato Formation.

Derivation of name: named after its smaller size compared to the type species.

Diagnosis and description (Figure 11.9f): hexagenitid larva with enlarged seventh gill of trapezoid shape, differs from type species by body length of 15.5 mm, length of terminal filaments about 6 mm (comparatively shorter than in type species). General body shape is drop-like (maximum width of body, 6.1 mm). Length of gill VII is about 2.4 mm. Otherwise it is very similar to the type species, including the larval wing pads that are medially partially fused (Figure 11.9f). Adults are still unknown.

Oligoneuriidae

Colocrus McCafferty, 1990 (see Figure 11.4b)

Cratoligoneuriella Martins-Neto, 1996a Syn. nov.

Diagnosis: larva: abdomen with pronounced posterolateral projections. Paracercus significantly shorter than cerci. Siphlonuroid setation of terminal filaments: cerci at their inner sides and terminal filum with swimming hairs. Abdomen with leaf-like tracheal gills preserved on segments I–VI (I–III in holotype AMNH 43483, III–V in SMNS larva 66538, I–VI in SMNS larva 66619). Gill I with dorsal insertion (McCafferty, 1990). Ventral gill tufts are not present or not preserved. Abdominal segments with posterolateral projections are present on segments IV–IX. Fore femora and fore tibiae on inner sides are equipped with a row of filtering setae (Plate 8d). Head capsule is broadened; the labium with enlarged and basally fused glossae forms a labial plate (Figure 11.9h; Plate 8e).

Alate stages: forewing with crossveins in costal field only. Gemination of Sc and R₁, R₄₊₅ and MA₁, MA₂ and MP₁, MP₂ and CuA. Series of veins running from CuA to posterior wing margin. Intercalaries absent except IMP.

Comments: Martins-Neto (1996a) established a new genus, *Cratoligoneuriella*, based on putative differences of the femoral base of the larval filter hairs. He claimed the presence of a ‘sclerotic process in the middle length of the fore femur’ of *Cratoligoneuriella* without further explanation or figure. Judging from the picture he gives, there is no difference between the detailed structure and origin of the filtering hairs in *Cratoligoneuriella* and *Colocrus*. However, the row of filtering hairs is sometimes detached and dislocated in some specimens (Figure 11.9g; Plate 8d). It is likely that Martins-Neto’s observation can be traced back to this artefact. I regard *Cratoligoneuriella leonardii* as a synonym of *Colocrus*.

Martins-Neto (2002b), in his unpublished doctoral thesis, described another species, *Cratoligoneuriella ninae*, and listed this species in a recent publication (Martins-Neto, 2005a). As the description in this doctoral thesis was never properly published, *Cratoligoneuriella ninae* has to be considered as a *nomen nudum*. This thesis was not available to me, so that I cannot comment on the description given there.

Colocrus indivicum McCafferty, 1990

Cratoligoneuriella leonardii (Martins-Neto, 1996a) Syn. nov.

Cratoligoneuriella ninae (Martins-Neto, 2005b) *nomen nudum*

Material: larval holotype AMNH 43484; adult paratype AMNH 43499 (Figure 11.4b); larvae SMNS 66538 (Figure 11.9g; Plate 8d) and SMNS 66619 (Plate 8e); larva RGMN-T005 coll. Martins-Neto; larvae MB.I.2025 (Figure 11.9h) and MB.I.2027 at MNB.

Diagnosis: larva (Figures 11.9g and h; Plate 8d and e): body length up to 15 mm excluding terminal filaments. Length of cerci up to 7 mm. Fore femora short and broad, middle and hind legs longer and slender. Filtering hairs on forelegs.

Alate stage (Figure 11.4b): forewing length 11 mm. Body length 13.2 mm. Wing venation see generic diagnosis, above.

Comment: McCafferty (1990) assigned an adult specimen (Figure 11.4b) of an oligoneuriid with a body length of 13.2 mm to *Colocrus indivicum*. Judging from the size of the holotype larva, it is most likely that it is indeed the alate stage of *C. indivicum*. Figure 11.9h and Plate 8e show in excellent preservation the labial plate that is characteristic for the larvae of all Oligoneuriidae. The filter apparatus is very well visible in specimen SMNS 66538 (Figure 11.9g; Plate 8d).

Colocrus? *magnus* sp. nov.

Material: larval holotype SMNS 66624 (Figure 11.9i). Putative adults are not rare in the collections; for example, specimen SMNS 66623 (Figures 11.6a and b), which is a fossil oligoneuriid of corresponding size, but the wing venation (Figures 11.6a and b) differs significantly from the adults attributed to *Colocrus indivicum*, so that the generic attribution is only very preliminary.

Type locality: Chapada do Araripe, vicinity of Nova Olinda, southern Ceará, north-east Brazil.

Type horizon: Lower Cretaceous, Upper Aptian, Nova Olinda Member of the Crato Formation.

Derivation of name: named after its large body size compared to the other species of the genus.

Diagnosis and description (Figure 11.9i): body length 22.5 mm without terminal filaments. Length of cerci 12 mm; length of terminal filament 9 mm. Femora short and broad.

Ephemeroidea (*sensu* McCafferty, 1991)

There are currently eight different species in eight different genera of Ephemeroidea described from Crato by McCafferty (1990) and Martins-Neto and Caldas (1990). While McCafferty (1990) described only alate stages, the contribution of Martins-Neto and Caldas (1990) only deals with larval specimens. A definite attribution of all larval and imaginal specimens to each other cannot be made at present.

Moreover, the larval specimens are not adequately described, and judging from the figures presented there, the contribution of Martins-Neto and Caldas rather deals with three specimens of a single species than three different genera and species from three different families. However, as I do not have the type material at hand, I here still refrain from synonymizing the three larval specimens described by Martins-Neto and Caldas (1990). The material and diagnoses given by the authors are merely listed.

Four undetermined larval specimens (two specimens SMNS 66630 and SMNS 66634, as well as two specimens without number in collections MURJ and MSF, respectively) and an undetermined adult specimen (MSF without number) of Ephemeroidea are featured in Figures 11.5a–c and 11.6h and i and Plate 8f.

Potamanthidae?

Olindinella gracilis Martins-Neto and Caldas, 1990

Material: larval holotype AMA-I-028 (Figure 11.4c), coll. Federal University of Ceará.

Diagnosis: burrowing larva with rostrum, long mandibular tusk, pubescent, robust tibia, paracercus longer than cercus.

Comment: validity and status doubtful, as discussed above.

Euthyplociidae

Pristiplocia rupestris McCafferty, 1990

Material: adult holotype AMNH 44308 (Figure 11.4d); adult SMNS 66541 (Figures 11.6c and d); various adult specimens in coll. MSF; larvae SMNS 66539 (Figure 11.6e) and SMNS 66621 (Figure 11.6f; Plates 8g and h) (probably two different species).

Diagnosis: alate stage (Figures 11.4d and 11.6c and d): body length 13–14 mm. Forelegs developed, about 4 mm long. Length of forewing 13 mm. Extensive crossvenation throughout wing. RS fork at one-quarter distance from base, MA fork in basal third of wing, MP₂ and CuA strongly decurved posteriorly at base. Cubital field with sigmoid veinlets from CuA to hind margin. A₁ not forked.

Larva (specimen SMNS 66621; Figure 11.6f; Plates 8g and h): body length up to 12 mm excluding terminal filaments (Figure 11.6f). Length of cerci and paracercus up to 2 mm. Abdominal segments II(?)–VII with fringed, doubled, lateral tracheal gills (Plate 8g). Femora flat and broad. Tibial spur not visible. Long, convex mandibular tusks up to 3 mm length, with numerous hairs (Plate 8h).

Comment: the very characteristic larval mandibular tusks equipped with long hairs unambiguously determine the larva as belonging to the Euthylociidae. As there have been only adult specimens of *P. rupestris* found so far, and because there are probably at least two species (Figures 11.6e and f) of euthyplociid larvae (compare the different length and shape of mandibles) from this locality, a definite attribution of one of these two types to the present species is not possible.

Ephemeridae

Australiephemera revelata McCafferty, 1990

Material: adult holotype AMNH 44300 (Figure 11.4e); adult paratype AMNH 44310 (Figure 11.4e).

Diagnosis: ephemeroïd forewing, MA forked at midlength, distinct distal arch of CuP, A₁ nearly straight, no cubital intercalaries, hind wing with costal angulation, RS shorter than R₂ and R₄₊₅.

Microephemera neotropica McCafferty, 1990

Material: adult holotype AMNH 43303 (Figure 11.4f).

Diagnosis: ephemeroïd forewing, MA forked in basal half of wing, moderate distal arch of CuP, A₁ not forked.

Cratonympha microcelata Martins-Neto & Caldas, 1990

Material: larval holotype USP GP/1T-1677 (Figure 11.4g).

Diagnosis: burrowing nymphs with rounded head, mandible long and smooth, foreleg with expanded tibia, apically with at least two spines.

Comment: validity and status doubtful, as discussed above.

Polymitarcyidae?

Caririnympha mandibulata Martins-Neto & Caldas, 1990

Material: larval holotype GP/1T-1678 coll. University of São Paulo.

Diagnosis: burrowing nymphs with trapezoid head, without rostrum, mandibular tusk short, foreleg with femur and tibia of same length.

Comment: validity and status doubtful, as discussed above.

Baetiscidae

Protobaetisca gen. nov.

Type species: *Protobaetisca bechlyi* sp. nov., by present designation.

Derivation of name: compound noun made up of the Greek prefix *protos* (first) and the Recent genus *Baetisca*.

Diagnosis: as for type species, since it is monotypic.

Protobaetisca bechlyi sp. nov.

Material: larval holotype SMNS 66620 (Figure 11.6g).

Type locality: Chapada do Araripe, vicinity of Nova Olinda, southern Ceará, north-east Brazil.

Type horizon: Lower Cretaceous, Upper Aptian, Nova Olinda Member of the Crato Formation.

Derivation of name: named after my colleague Günter Bechly, curator of fossil insects and amber at the SMNS, who pointed me to fossil mayflies.

Diagnosis and description (Figure 11.6g): larva with stout, rotund body of ovoid shape. Length 8 mm without terminal filaments. Length of cerci 3 mm. Abdominal segments VI–IX with prominent posterolateral projections, without median spines. Lateral margins of abdominal segments I–V bent up. Abdominal segment VI slightly enlarged, longest abdominal segment, with prominent longitudinal crest most probably representing the caudal closure of a gill chamber. Mesonotum posteriorly extending up to abdominal segment VI, forming a gill chamber ('carapace'). Anterior part of gill chamber torn and flaked off. It cannot be decided with certainty if the pronotal part of the gill chamber is preserved in its entity or partly broken off together with the anterior part of the mesonotum). Carapace without dorsal spines, lateral parts of carapace not preserved. Length of preserved part of foreleg 4 mm, length of femur 2 mm, length of tibia 0.8 mm. Head with two frontal carinae. Antennae not preserved.

Comment: apparently, this is the fourth record of a fossil stem group representative of the Prosopistomatoidea. Staniczek and Bechly (2002) described *Balticobaetisca velteni* from Eocene Baltic amber extending the distribution of the

Baetiscidae to the Palaearctic. Jell and Duncan (1986) described a specimen from the Lower Cretaceous as Siphlonuridae? gen. nov. It has the general shape of a baetiscid mayfly with a mesonotum that extends posteriorly to abdominal segment VI. The pronotum is separated from the mesonotum. The authors mentioned its similarities to Baetiscidae but refrained from assigning the fossil to this family because its Recent distribution is Nearctic. Nevertheless this fossil either belongs to the stem group of Baetiscidae or Prosopistomatoidea. Sinitshenkova (2000) described *Cretomitarcys luzzi* from the Upper Cretaceous of New Jersey. She assigned the adult male specimen to the †Cretomitarcyinae, a new subfamily of the Polymitarcyidae. McCafferty (2004) realized its affinities and transferred the fossil to the Prosopistomatoidea (= Carapacea *sensu* McCafferty = Posteritorna *sensu* Kluge). He placed it in a new family, the †Cretomitarcyidae. Judging from the figures given in Sinitshenkova (2000), *C. luzzi* indeed clearly belongs to the Prosopistomatoidea: the wing tornus is situated posterior to A₁. The rounded hind wing with multiple intercalaries points to a phylogenetic position in the stemline of Baetiscidae. *C. luzzi* is plesiomorphic in retaining gonopods with several segments and an entirely separated basitarsus of the foreleg. However, I see no reason in placing this species in a separate family.

Protobaetisca bechlyi sp. nov. is the first record of a fossil Baetiscidae from the Crato Formation. The frontal head projections confirm its placement at least in the stem group of Baetiscidae. If the pronotum was separated a placement in the stem group of Prosopistomatoidea cannot be excluded.

The alleged fossil Baetiscidae *Caririephemera marquesi* (Zamboni, 2001) definitely does not belong to the Baetiscidae (see below).

Other records of unnamed Ephemeroptera

- Siphlonuridae (?) sp. 1 McCafferty (1990: 30, figure 11; AMNH 44306, adult)
- Siphlonuridae (?) sp. 2 McCafferty (1990: 30, figure 12; AMNH 44313, adult)
- Siphlonuridae (?) sp. 3 McCafferty (1990: 31, figure 13; AMNH 43477, adult)
- Oligoneuriidae gen. et sp. indet. Martins-Neto (1996a: 188, figure 6c; adult)
- Ephemeroidea *incertae sedis* (Figures 11.5a and b and 11.6h and i; Plate 8f); larval specimens SMNS 66630, SMNS 66634, without number in collections MURJ and MSF)
- Ephemeroidea *incertae sedis* (Figure 11.5c; adult specimen, MSF without number)
- Ephemeroidea sp. 1 McCafferty (1990: 42, figure 30; AMNH 44311, adult)
- Ephemeroidea sp. 2 McCafferty (1990: 42–43, figure 31; AMNH 43480, adult)
- Leptophlebiidae (?) sp. 1 McCafferty (1990: 43, figure 32; AMNH 43474, adult)
- *Comment:* this taxon is relatively frequently found. Other specimens (e.g. SMNS 66639, Figure 11.5d) demonstrate that specimen AMNH 43474 is not a larva, but an adult without preserved wings. The other fossils generally have entangled or wizened wings (probably subimagines) and also only two terminal filaments (cerci).

- Leptophlebiidae (?) sp. 2 McCafferty (1990: 43–44; figure 33; AMNH 43476, adult)
- Leptophlebiidae (?) sp. 3 McCafferty (1990: 44, figure 34; AMNH 44312, adult)
- Family *incertae sedis* sp. 1 McCafferty (1990: 44; AMNH 43453, larva)
- Family *incertae sedis* sp. 2 McCafferty (1990: 44–45; AMNH 43423, larva)
- Family *incertae sedis* Grimaldi and Engel (2005: 166, figure 6.12; AMNH without number; small alate stage with two extremely long cerci but without paracercus)
- Family *incertae sedis* Bechly *et al.* (2001a: figure 30)
- Familia *incertae sedis* (Figures 11.5e and f); larval specimens SMNS 66622, SMNS 66625 and MSF Z2)
- *Comment:* these strange mayfly larvae (body length about 23 mm) have an absolutely unique habitus with broadened and flattened fore and hind femora, but slender mid femora.

Ephemeroptera incertae sedis

Costalimella nordestina Martins-Neto, 1996a (Figure 11.8a)

Costalimella zuechii Zamboni, 2001

Comment: these two species have each been described from a single adult specimen. Possibly both specimens resemble small species of †Hexagenitidae. In the forewing, the MA branch is located in the apical fourth of the wing, and in *C. zuechii* the branching of CuA is reported.

Insecta incertae sedis

Caririephemeria marquesi Zamboni, 2001

Comment: Zamboni, 2001 described an insect larva with eight visible abdominal segments and without abdominal gills as a mayfly larva. Terminal filaments are either not present or not preserved. The author tentatively places the fossil within the Baetiscidae, because ‘the lack of gills . . . occurs only in Baetiscidae’. In fact, the gills of the Baetiscidae lie under a gill chamber formed by the enlarged mesonotum. This fossil has no such gill chamber and nor does it exhibit any character that could identify it as a mayfly nymph, even though it might well represent a poorly preserved specimen of *Protoligoneuria limai*.

11.5 Odonata: damselflies and dragonflies

Günter Bechly

The order Odonata includes three Recent suborders (Zygoptera, ‘Anisozygoptera’ – Epiophlebiidae, and Anisoptera) with a total of 635 Recent genera and 5,538 described species. Odonates are relatively large insects and well known for their beautiful colours, their swift flight and the curious mating in the wheel-position.

Odonates have bristle-like antennae, biting mouthparts and large compound eyes. The thoracic segments have a distinct skew, so that their spiny legs are tilted

belong to the same genus. A careful revision by a specialist on fossil Trichoptera would be very useful. Furthermore, there are several putative new genera and species to be described, such as specimens nos SMNS 66282 (Figure 11.87d), SMNS 66287 (Figure 11.88a) and SMNS 66568 (Figure 11.88b), which are clearly distinct from any described species.

Crato Lepidoptera: moths and butterflies

Lepidoptera are extremely rare in the Crato Formation and only members of the microlepidopteran grade occur. Martins-Neto and Vulcano (1989b) and Martins-Neto (1999, 2001b) described the following five taxa.

- *Parasabatinca caldasae*: Micropterygidae; body length 3.5–5 mm; forewing length 3–3.5 mm; R_2 and R_3 ending near apex. Besides the holotype, a single specimen (SMNS 66279) has been identified as this species (Figure 11.87c).
- *Undopterix caririensis*: Undopterygidae; hind wing length 3.1 mm; M_1+M_2 fork slightly posterior to R_4+R_5 fork; CuA ending at anal margin near apex.
- *Gracilepterix pulchra*: in familia incertae sedis; similar to *Undopterix*; body length and forewing length about 3 mm; hind wing length 2.3 mm; R_1 forking anteriorly of R_2+R_3 and R_4+R_5 forks in forewings; Sc and R_1 unbranched in hind wings. One specimen of this species (SMNS 66277) has been identified (Figure 11.87d). This fossil butterfly was incorrectly listed as *Gracilepteryx* (*sic*) by Martins-Neto (2005b).
- *Xena nana* Martins-Neto, 1999b: Eolepidopterigidae. This taxon seems to be similar to *Eolepidopterix* and is said to be described by ‘Martins-Neto, 2000’ according to Martins-Neto (2005b), which seems to be a *lapsus* and should be ‘Martins-Neto, 1999b’.
- *Psamateia calipsa* Martins Neto, 2002: the present author could not locate the publication with this description, which is mentioned by Martins-Neto (2005b).

All the above species belong to the most primitive and most basal grade of Lepidoptera, and thus not within the Glossata, although Martins-Neto (2005b: 480) attributed all five species to the Eolepidopterigidae, but without explication. However, Grimaldi and Engel (2005: 562, figure 13.16) figured an unnamed primitive moth (AMNH SF 46441) with a piercing oviscapt similar to the Recent families Eriocraniidae and Acanthopteroctetidae that are indeed basal Glossata. This fossil could be the most derived Lower Cretaceous lepidopteran yet discovered, if it should not turn out to be just a small trichopteran like *Cratorella media*.

References

- Achtelik, M. 1967. Ueber die Anatomie des Kopfes von *Raphidia flavipes* Stein und die Verwandtschaftsbeziehungen der Raphidiidae zu den Megaloptera. *Zoologische Jahrbücher, Abteilung für Anatomie und Ontogenie der Tiere* **84**: 249–312.

- Afzelius, B. A. and Dallai, R. 1988. Spermatozoa of Megaloptera and Raphidioptera (Insecta, Neuropteroidea). *Journal of Ultrastructural and Molecular Structure Research* **101**: 185–191.
- Alexander, B. A. 1992. An exploratory analysis of cladistic relationships within the superfamily Apoidea, with special reference to sphecid wasps (Hymenoptera). *Journal of Hymenoptera Research* **1**: 25–61.
- Alexeev, A. V. 1994. Jurassic and Lower Cretaceous Buprestidae (Coleoptera) from Eurasia. *Paleontological Journal* **27**(I A): 9–34.
- 1996. New taxa of metallic wood-boring beetles (Coleoptera, Buprestidae) from the Mesozoic of Russia, Kazakhstan, and Mongolia. *Paleontological Journal* **30**: 310–117.
- 2000. On Mesozoic Buprestids (Coleoptera: Buprestidae) from Russia, Kazakhstan, and Mongolia. *Paleontological Journal* **34** (suppl. 3): 323–326.
- Andersen, N. M. 1998. Water striders from the Paleogene of Denmark with a review of the fossil record and evolution of semiaquatic bugs (Hemiptera: Gerromorpha). *Det Kongelige Danske Videnskabernes Selskab, Biologiske Skrifter* **50**: 1–152.
- and Weir, T. A. 2004. Australian water bugs. Their biology and identification (Hemiptera-Heteroptera, Gerromorpha and Nepomorpha). *Entomonograph* **14**: 1–344.
- Andersen, S. 2001. Silky lacewings (Neuroptera: Psychopsidae) from the Eocene–Palaeocene transition of Denmark with a review of the fossil record and comments on phylogeny and zoogeography. *Insect Systematics and Evolution* **32**: 419–438.
- Anderson, R. S. 2005. New Oxycoryninae from Central and South America: phylogenetic and biogeographical implications (Coleoptera: Belidae). *Systematic Entomology* **30**: 644–652.
- Ansorge, J. and Mostovski, M. B. 2000. Redescription of *Prohirmoneura jurassica* Handlirsch 1906 (Diptera: Nemestrinidae) from the Lower Tithonian lithographic limestone of Eichstätt (Bavaria). *Neues Jahrbuch für Geologie und Paläontologie Monatshefte* **2000**: 235–243.
- Anufriev, G. A. and Emeljanov, A. F. 1988. Tsikadovye (Auchenorrhyncha, Cicadinea). [Suborder Cicadinea (Auchenorrhyncha)], pp. 12–495. In Lehr, A. P. (ed), *Oprelelitel' nasekomykh Dal'nego Vostoka SSSR*. St Petersburg: Nauka.
- Arnol'di, L. V., Zherikhin, V. V., Nikritin, L. M. and Ponomarenko, A. G. 1992. *Mesozoic Coleoptera*. Washington DC: Smithsonian Institution Libraries and The National Science Foundation.
- Asche, M. 1988. Preliminary thoughts on the phylogeny of Fulgoromorpha (Homoptera, Auchenorrhyncha). *Proceedings of the 6th Auchenorrhyncha Meeting*, Turin, Italy, 7–11 September 1987, pp. 47–53.
- Aspöck, U. and Aspöck, H. 2004. Two significant new snakeflies from Baltic amber, with discussion on autapomorphies of the order and its included taxa (Raphidioptera). *Systematic Entomology* **29**: 11–19.
- Azar, D. and Nel, A. 2002. New Cretaceous psychodid flies from Lebanese amber and Santana Formation (Chapada do Araripe, Brazil) (Diptera). *Annales de la Societe Entomologique de France (n.s.)* **38**: 253–262.
- Baudoin, R. 1980. Sur les Gerris des miroirs d'eau actuels et les Chresmoda des lagunes post-récifales portlandiennes de Solnhofen. *Annales des Sciences Naturelles / Zoologie* **2**: 111–116.
- Bechly, G. 1996. Review: SANTANA 1.0 (1996): CD-ROM by digital-fossil. *Petalura* **2**: <http://www.bernstein.naturkundemuseum-bw.de/odonata/pet2p03.htm>.

- 1997a. Die Libellen (und einige andere fossile Insekten) aus der Unterkreide von Brasilien (Santana). 4. *Fachgespräch "Fossile Insekten"*, Clausthal-Zellerfeld, 28–29 June 1997, Abstracts.
 - 1997b. Dragonflies from the Lower Cretaceous of Brazil. *Meganeura* **1**: 27–28.
 - 1997c. Dragonflies from the Lower Cretaceous of Brazil. *Inclusion Wrostek* **27**: 9.
 - 1998a. Santana – Schatzkammer fossiler Insekten. *Fossilien* **2/98**: 95–99.
 - 1998b. Santana – Forschungsgeschichte und Fauna. *Fossilien* **3/98**: 148–156.
 - 1998c. New fossil dragonflies from the Lower Cretaceous Crato Formation of north-east Brazil (Insecta: Odonata). *Stuttgarter Beiträge zur Naturkunde, Serie B* **264**: 1–66.
 - 1998d. [Santana – Eldorado of Mesozoic fossils]. *Catalogue of the Taipei Natural History Show* **1998**: 148–156.
 - 1999a *Phylogeny and Systematics of Fossil Dragonflies (Insecta: Odonatoptera) with Special Reference to some Mesozoic Outcrops*. Published PhD thesis, Eberhard-Karls-Universität Tübingen.
 - 1999b. Notices. *Meganeura* **4**: 9.
 - 2000. Two new fossil dragonfly species (Insecta: Odonata: Anisoptera: Araripegomphidae and Lindeniiidae) from the Crato Limestone (Lower Cretaceous, Brazil). *Stuttgarter Beiträge zur Naturkunde, Serie B* **296**: 1–16.
 - 2002. Phylogenetic systematics of Odonata. In Schorr, M. and Lindeboom, M. (eds), *Fundamentals of Odonatological Research*. ODOLITonCD 1.0 (ISSN 1438–034x). www.dragonflyresearch.de/.
 - 2003. The phylogenetic relationships of the three extant suborders of Odonata. *Entomologische Abhandlungen* **61**: 127–128.
 - 2005. A re-description of "Stenophlebia" casta (Insecta: Odonata: Parastenophlebiidae n. fam.) from the Upper Jurassic Solnhofen Limestone in Germany. *Stuttgarter Beiträge zur Naturkunde, Serie B* **359**: 1–12.
 - and Ueda, K. 2002. The first fossil record and first New World record for the dragonfly clade Chlorogomphida (Insecta: Odonata: Anisoptera: Araripechlorogomphidae n. fam.) from the Crato Limestone (Lower Cretaceous, Brazil). *Stuttgarter Beiträge zur Naturkunde, Serie B* **328**: 1–11.
 - , Nel, A. Martínez-Delclòs, X. and Fleck, G. 1998. Four new dragonfly species from the Upper Jurassic of Germany and the Lower Cretaceous of Mongolia (Anisoptera: Hemeroscopidae, Sonidae, and Proterogomphidae fam. nov.). *Odonatologica* **27**: 149–187.
 - , Haas, F., Schwaller, W., Schmalfuss, H. and Schmid, U. 2001a. Ur-Geziefer – Die faszinierende Evolution der Insekten. *Stuttgarter Beiträge zur Naturkunde, Serie C* **49**: 1–96.
 - , Brauckmann, C., Zessin, W. and Gröning, E. 2001b. New results concerning the morphology of the most ancient dragonflies (Insecta: Odonatoptera) from the Namurian of Hagen-Vorhalle (Germany). *Journal of Zoological Systematics and Evolutionary Research* **39**: 209–226.
 - , Nel, A. Martínez-Delclòs, X. et al. 2001c. A revision and phylogenetic study of Mesozoic Aeshnoptera, with description of several new families, genera and species (Insecta: Odonata: Anisoptera). *Neue paläontologische Abhandlungen* **4**: 1–219.
 - , Dietl, G. and Schweigert, G. 2003. A new species of *Stenophlebia* (Insecta: Odonata: Stenophlebiidae) from the Nusplingen Lithographic Limestone (Upper Jurassic, SW Germany). *Stuttgarter Beiträge zur Naturkunde, Serie B* **338**: 1–10.
- Becker-Migdisova, E. E. 1946. Ocherki po sravnitel'noi morfologii sovremennykh i permskikh Homoptera. Chast' I. *Izvestia AN SSSR, Biology Series* **6**: 741–766.

- 1947. *Cicadoprosbole sogutensis* gen. et sp. nov. – perekhodnaya forma mezhdu permeskimi Prosbolidae i sovremennymi Cicadidae. *Doklady Akademii Nauk SSSR* 5: 445–448.
- 1949. Mezozojskie Homoptera Srednei Azii. *Trudy Palaeontologicheskogo Instituta Nauk* 22: 1–68.
- 1962a. Nekotorye novye poluzhestokrylye i senoedy. *Palontologicheskiy Zhurnal* 1: 89–104.
- 1962b. Otryad Homoptera. Ravnokrylye, pp. 162–208. In Rohdendorf, B. B. (ed.), *Osnovy paleontologii. Chlenistonogie trakheinye i khelitserovye*. Moscow: Academy of Sciences USSR.
- and Popov, Yu., A. 1963. Some representatives of Heteroptera from the Jurassic of Karatau. *Palaeontological Journal of the Academy of Science USSR* 2: 74–82.
- Bedford, G. O. 1978. The biology and ecology of the Phasmatodea. *Annual Review of Entomology* 23: 125–149.
- Beier, M. 1968. Mantodea (Fangheuschrecken). *Handbuch der Zoologie*, vol. IV, Band Arthropoda, 2 Hälfte: Insecta. Zweite Auflage: 1–47.
- Belayeva, N. V. 2002. Order Termitida, pp. 270–273. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
- Bennett, D. J. and Engel, M. S. 2005. A primitive sapygid wasp in Burmese amber (Hymenoptera: Sapygidae). *Acta Zoologica Cracoviensia* 48B(1–2): 1–9.
- Béthoux, O. and Nel, A. 2002. Venation pattern and revision of Orthoptera sensu nov. and sister groups. Phylogeny of Palaeozoic and Mesozoic Orthoptera sensu nov. *Zootaxa* 96: 1–88.
- Nel, A., Lapeyrie, J., Gand, G. and Galtier, J. 2002. *Raphogla rubra* gen. n., sp. n., the oldest representative of the clade of modern Ensifera (Orthoptera: Tettigoniidea, Gryllidea). *European Journal of Entomology* 99: 111–116.
- Beutel, R. G. 1995. The Adephaga (Coleoptera): phylogeny and evolutionary history, pp. 173–217. In Pakaluk, J. and Ślipiński, S. A. (eds), *Biology, Phylogeny, and Classification of Coleoptera*, vol. 1. Warszawa: Muzeum I Instytut Zoologii PAN.
- Bischoff, H. 1927. *Biologie der Hymenopteren, Eine Naturgeschichte der Hautflügler*. Biologische Studienbücher. Berlin: J. Springer.
- Bitsch, C. and Bitsch, J. 2000. The phylogenetic interrelationships of the higher taxa of apterygote insects. *Zoologica Scripta* 29: 131–156.
- Blagoderov, V., Grimaldi, D. A. and Fraser, N. C. 2007. How time flies for flies: diverse Diptera from the Triassic of Virginia and early radiation of the order. *American Museum Novitates* (in press).
- Bode, A. 1953. Die Insektenfauna der Ostniedersächsischen oberen Liás. *Palaeoentomographica A* 103: 1–406.
- Bohart, R. M. and Menke, M. S. 1976. *Sphecid wasps of the world. A generic revision*. Los Angeles: University of California Press, Berkeley.
- Bohn, H. 2003a. Ordnung Blattoptera, Schaben, pp. 197–223. In Dathe, H. H. (ed), *Lehrbuch der Speziellen Zoologie, Band 1, Teil 5 (Insecta)*. Heidelberg: Spektrum.
- 2003b. Ordnung Isoptera, Termiten, pp. 223–250. In Dathe, H. H. (ed.), *Lehrbuch der Speziellen Zoologie, Band 1, Teil 5 (Insecta)*. Heidelberg: Spektrum.
- Bordy, E. M., Bumby, A. J., Catuneanu, O. and Eriksson, P. G. 2004. Advanced early Jurassic termite (Insecta: Isoptera) nests: evidence from the Clarens Formation in the Tuli Basin, southern Africa. *Palaios* 19: 68–78.
- Börner, C. 1909. Neue Homologien zwischen Crustaceen und Hexapoden. Die Beißmandibel der Insekten und ihre phylogenetische Bedeutung. Archi- und Metapterygota. *Zoologischer Anzeiger* 34: 100–125.

- Boudreaux, H. B. 1979. *Arthropod Phylogeny with Special Reference to Insects*. New York: Wiley.
- Boukary, I. B., Tourneur, J. C. and Gingras, J. 1996. Life cycle of *Forficula senegalensis* Serv. (Dermoptera: Forficulidae) and its relationship to the development of bulrush millet in Sundanese-Sahelian zone of Niger. *Canadian Entomologist* **128**: 831–838.
- Boulard, M. and Nel, A. 1990. Sur deux cigales fossiles des terrains tertiaires de la France [Homoptera, Cicadoidea]. *Revue Française d'Entomologie (n.s.)* **12**: 37–45.
- Bourgoin, Th. 1993. Cladistic analysis of the Meenopliidae–Kinnaridae genera: the Kinnaridae, a paraphyletic family (Hemiptera, Fulgoromorpha). In Drosopoulos, S., Petrakis, P. V., Claridge, M. F. and de Vrijer, P. W. F. (eds), *Proceedings of the 8th Auchenorrhyncha Congress*, Delphi, Greece, 9–13 August 1993: 22–24.
- and Campbell, B. C. 2002. Inferring a phylogeny for Hemiptera: falling into the ‘autapomorphic trap’, pp. 67–82. In Holzinger, W. (ed.), *Zikaden – Leafhoppers, Planthoppers and Cicadas (Insecta: Hemiptera: Auchenorrhyncha)*. Denisia **04**, Zugleich Kataloge, Neue Folge Nr. 176, Linz: OÖ. Landesmuseums.
- Steffen-Campbell, J. D. and Campbell, B. C. 1997. Molecular phylogeny of Fulgoromorpha (Insecta, Hemiptera, Archaeorrhyncha). The enigmatic Tettigometridae: evolutionary affiliations and historical biogeography. *Cladistics* **17**: 207–224.
- Bradley, C. 1955. Notes on the synonymy, distribution and affinities of the subfamily Fedtschenkiinae of the Sapygidae (Hymenoptera). *Entomological News* **66**: 230–233.
- Brandao, C. R. F., Martins-Neto, R. G. and Vulcano, M. A. 1989. The earliest known fossil ant (First Southern Hemisphere mesozoic record) (Hymenoptera: Formicidae: Myrmeciinae). *Psyche* **96**: 195–208.
- Brito, I. M. 1984. Nota preliminar sobre os insetos da Formação Santana, Cretáceo inferior da Chapada do Araripe. *XXXIII Congresso Brasileiro de Geologia, Rio de Janeiro*: 530–535.
- 1987. Nota preliminar sobre uma nova efêmera do Cretáceo do Ceará (Insecta Ephemeroptera). *Anais do X Congresso Brasileiro de Paleontologia* **2**: 593–597.
- Brogniart, C. 1893. *Recherches pour servir à l'Histoire des Insectes Fossiles des Temps Primaires, Précédées d'une Étude sur la Nervation des Ailes des Insects*. Thèse présentée à la Faculté des Sciences de Paris, no. 821; Paris.
- Brothers, D. J. 1975. Phylogeny and classification of the aculeate Hymenoptera, with special reference to Mutillidae. *University of Kansas Science Bulletin* **50**: 483–648.
- Burckhardt, D. and Agosti, D. 1991. New records of South American Peloridiidae (Homoptera: Coleorrhyncha). *Revista Chilena Entomología* **19**: 71–75.
- and Cekalovic, T. K. 2002. An anomalous moss-bug from Southern Chile and notes on *Pantinia darwini* (Hemiptera, Coleorrhyncha, Peloridiidae). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft / Bulletin de la Société entomologique Suisse* **75**: 57–59.
- Burnham, L. 1978. Survey of Social Insects in the Fossil Record. *Psyche* **85**: 85–133.
- Carle, F. L. 1982. The wing vein homologies and phylogeny of the Odonata: a continuing debate. *S.I.O. Rapid Communications* **4**: 1–66.
- 1995. Evolution, taxonomy, and biogeography of ancient Gondwanian Libelluloids, with comments on anisopterid evolution and phylogenetic systematics (Anisoptera: Libelluloidea). *Odonatologica* **24**: 383–424.
- and Wighton, D. C. 1990. Odonata, pp. 51–68. In Grimaldi, D. A. (ed.), *Insects from the Santana Formation, Lower Cretaceous, of Brazil*. *Bulletin of the American Museum of Natural History*, vol. 195.

- Carpenter, F. M. 1933. The Lower Permian Insects of Kansas. Part 6. *Proceedings of the American Academy of Arts and Sciences* 68: 411–503.
- 1979. The Lower Permian insects from Oklahoma. Part 2. Orders Ephemeroptera and Palaeodictyoptera. *Psyche* 86: 261–290.
- 1992. Superclass Hexapoda. In Kaesler, R. L. (ed), *Treatise on Invertebrate Paleontology, R, Arthropoda 4, 3/4*. Boulder and Lawrence: Geological Society of America and University of Kansas.
- Carpenter, J. M. 1982. The phylogenetic relationships and natural classification of the Vespoidea (Hymenoptera). *Systematic Entomology* 7: 11–38.
- 1987. Phylogenetic relationships and classification of the Vespinae (Hymenoptera: Vespidae). *Systematic Entomology* 12: 423–431.
- and Rasnitsyn, A. P. 1990. Mesozoic Vespidae. *Psyche* 97(1–2): 1–20.
- Carver, M., Gross, G. F. and Woodward, T. E. 1992. Hemiptera, pp. 429–509. In Naumann, I. D. (ed), *The Insects of Australia*. Melbourne: CSIRO and Melbourne University Press.
- Cassola, F. and Werner, K. 2004. A fossil tiger beetle specimen from the Brazilian Mesozoic: *Oxycheilopsis cretacicus* n. gen., n. sp. *Mitteilungen der Münchener entomologischen Gesellschaft* 94: 75–81.
- Chadwick, M. A., Hunter, H., Feminella, J. W. and Henry, R. P. 2002. Salt and water balance in *Hexagenia limbata* (Ephemeroptera: Ephemeridae) when exposed to brackish water. *Florida Entomologist* 85: 650–651.
- Chen, S. and Tian, C.-c. 1973. A new family of Coleoptera from the Lower Cretaceous of Kansu. *Acta Entomologica Sinica* 16: 169–179.
- Chin, K. and Gill, B. D. 1996. Dinosaurs, dung beetles, and conifers: participants in a Cretaceous food web. *Palaios* 11: 280–285.
- Cockerell, T. D. A. 1913. The first fossil Mydaid fly. *The Entomologist* 46: 207–208.
- 1917. Insects in Burmese amber. *Annals of the Entomological Society of America* 10: 323–329.
- Colgan, D. J., Cassis, G. and Beacham, E. 2003. Setting the molecular phylogenetic framework for the Dermaptera. *Insect Systematics and Evolution* 34: 60–80.
- Compton, S. G. and Ware, A. B. 1991. Ants disperse the elaiosome-bearing eggs of an African stick insect. *Psyche* 98: 207–213.
- Crampton, G. C. 1938. The interrelationships and lines of descent of living insects. *Psyche* 45: 165–181.
- Crowson, R. A. 1981. *The Biology of Coleoptera*. New York: Academic Press.
- Cryan, J. R. 2005. Molecular phylogeny of Cicadomorpha (Insecta: Hemiptera: Cicadoidea, Cercopoidea and Membracoidea): adding evidence to the controversy. *Systematic Entomology* 30: 563–574.
- Currie, D. C. and Grimaldi, D. 2000. A new blackfly (Diptera: Simuliidae) genus from mid-Cretaceous (Turonian) amber of New Jersey, pp. 473–485. In Grimaldi, D. (ed), *Studies on Fossils in Amber, with Particular Reference to the Cretaceous of New Jersey*. Leiden: Backhuys.
- da Costa-Lima, A. 1950. Ninfas de Efemerídeo fossil do Ceará. *Anais Academia, Brasileira, Ciências* 22: 419–420.
- Darling, D. C. and Sharkey, M. J. 1990. Order Hymenoptera, pp. 123–153. In Grimaldi, D. A. (ed.), *Insects from the Santana Formation, Lower Cretaceous, of Brazil. Bulletin of the American Museum of Natural History* 195.
- Day, M. C., Else, G. R. and Morgan, D. 1981. The most primitive Scoliidae (Hymenoptera). *Journal of Natural History* 15: 671–684.

- Deitz, L. L. and Dietrich, C. H. 1993. Superfamily Membracoidea (Homoptera: Auchenorrhyncha). I. Introduction and revised classification with new family-group taxa. *Systematic Entomology* **18**: 287–296.
- Demoulin, G. 1955. Sur une larve siphlonuridienne d'Ephémère fossile du Brésil. *Bulletin et Annales de la Société Royale d'Entomologie de Belgique* **91**: 270–271.
- 1971. Contribution à l'étude morphologique, systématique et phylogénique des Ephemeroptères Jurassiques (1). VI. L'aile postérieure des Hexagenites Scudder et les rapports Hexagenitidae-Chromarcyidae-Oligoneuriidae. *Bulletin* **47(29)**: 1–10.
- Desutter-Grandcolas, L. 2003. Phylogeny and the evolution of acoustic communication in extant Ensifera (Insecta, Orthoptera). *Zoologica Scripta* **32**: 252–261.
- Dietrich, C. H. 1999. The role of grasslands in the diversification of leafhoppers (Homoptera: Cicadellidae): a phylogenetic perspective, pp. 44–48. In Warwick, C. (ed.), *Proceedings of the Fifteenth North American Prairie Conference*. Bend, OR: Natural Areas Association.
- 2002. Evolution of Cicadomorpha (Insecta, Hemiptera), pp. 155–170. In Holzinger, W. (ed.), *Zikaden – Leafhoppers, Planthoppers and Cicadas (Insecta: Hemiptera: Auchenorrhyncha)*, Denisia, 4, Neue Folge Nr. 176, zugleich Kataloge des OÖ. Linz: Landesmuseums.
- 2005. Keys to the families of Cicadomorpha and subfamilies and tribes of Cicadellidae (Hemiptera: Auchenorrhyncha). *Florida Entomologist* **88**: 502–517.
- and Rakitov, R. A. 2002. Some remarkable new deltocephaline leafhoppers (Hemiptera: Cicadellidae: Deltocephalinae) from the Amazonian rainforest canopy. *Journal of the New York Entomological Society* **110**: 1–48.
- McKamey, S. H. and Dietz, L. L. 2001. Morphology-based phylogeny of the treehopper family Membracidae (Hemiptera: Cicadomorpha: Membracoidea). *Systematic Entomology* **26**: 213–239.
- Deitz, L. L. and Dietrich, C. H. 1993. Superfamily Membracoidea (Homoptera: Auchenorrhyncha). I. Introduction and revised classification with new family-group taxa. *Systematic Entomology* **18**: 287–296.
- Distant, W. L. 1883. Contribution to a proposed monograph of the homopterous family Cicadidae. *Proceedings of the Zoological Society of London* **1883**: 187–194.
- 1905. Rhynchotal notes XXXIII. *Annals and Magazine of Natural History* **7**: 1622–1635.
- Dolin, V. G. and Nel, A. 2002. Trois nouveaux Elateridae fossiles du Mésozoïque supérieur de Chine (Coleoptera). *Bulletin de la Société entomologique de France* **107**: 341–346.
- Ehrmann, R. 2002. *Mantodea. Gottesanbeterinnen der Welt*. Munich: Natur und Tier Verlag GmbH.
- Eichwald, E. 'd. 1864. Sur un terrain jurassique a poissons et Insectes d'eau douce de la Sibérie orientale. *Bulletin de Societe Geologique de France* **21**: 19–25.
- Emeljanov, A. F. 1991. K voprosu ob obeme i podrazdeleniyakh sem. Achlidiae (Homoptera, Cicadina). *Entomologicheskoe Obozrenie* **70**: 373–393. [Published in English as: Emel'yanov, A. F. 1992. Toward the problem of the limits and subdivisions of Achlidiae (Homoptera, Cicadina). *Entomological Review* **71**: 53–73].
- 1992. Opisanie trib podsem. Achilinae (Homoptera, Achilidae) i utochnenie ikh sostava. *Entomologicheskoe Obozrenie* **71**: 574–594. [Published in English as: Emel'yanov, A. F. 1993. Description of tribes of the subfamily Achilinae (Homoptera, Achilidae) and revision of their composition. *Entomological Review* **72**: 7–27].

- 1994 Pervaya iskopayemaya nakhodka semeistva Derbidae i pereopisanie paleogenovogo roda *Hooleya* Cockerell (Achilidae) (Insecta: Homoptera, Fulgoroidea). *Paleontologicheskii Zhurnal* **3**: 76–82. [Published in English as: Emeljanov, A. F. 1995. The first find of fossil Derbidae, and a redescription of Paleogene achilid *Hooleya* Cockerell (Insecta: Homoptera, Fulgoroidea). *Paleontological Journal* **28**: 92–101].
- 1999. Notes on the delimitation of families of the Issidae group with description of a new species of Caliscellidae belonging to a new genus and tribe (Homoptera, Fulgoroidea). *Zoosystematica Rossica* **8**: 61–72.
- 2002. Contribution to classification and phylogeny of the family Cixiidae (Hemiptera, Fulgoromorpha), pp. 103–112. In Holzinger, W. (ed), *Zikaden – Leafhoppers, Planthoppers and Cicadas (Insecta: Hemiptera: Auchenorrhyncha)* Denisia, 04, Zugleich Kataloge, Neue Folge Nr. 176, Linz: OÖ. Landesmuseums.
- Engel, M. S. 2001. A monograph of the Baltic amber bees and evolution of the Apoidea (Hymenoptera). *Bulletin of the American Museum of Natural History* **259**: 1–192.
- 2002. The smallest snakefly (Raphidioptera: Mesoraphidiidae): a new species in Cretaceous amber from Myanmar, with a catalogue of fossil snakelies. *American Museum Novitates* **3363**: 1–22.
- and Grimaldi, D. A. 2004. New light shed on the oldest insect. *Nature* **427**: 627–630.
- and Chatzimanolis, S. 2005. Early Cretaceous earwigs (Dermaptera) from the Santana Formation, Brazil. *Polskie Pismo Entomologiczne* **74**: 219–226.
- Esaki, T. 1949. The occurrence of the Mesozoic insect *Chresmoda* in the Far East. *Insecta Matsumarana* **17**: 4–5.
- Eskov, K. Yu. 1984. Dreif materikov i problema isoricheskoi biogeografi, pp. 24–92. In: *Faunogenet i filogenet*. Moscow: Nauka.
- and Golovatch, S. I. 1986. On the origin of Trans-Pacific disjunctions. *Zoologisches Jahrbuch für Systematik* **113**: 265–285.
- Evans, J. W. 1956. Palaeozoic and Mesozoic Hemiptera. *Australian Journal of Zoology* **4**: 165–258.
- 1981. A review of the present knowledge of the family Peloridiidae and new genera and new species from New Zealand and New Caledonia (Hemiptera: Insecta). *Records of the Australian Museum* **34**: 381–406.
- Evenhuis, N. L. 1994. *Catalogue of the Fossil Flies of the World (Insecta: Diptera)*. Leiden: Backhuys.
- Fanenbruck, M., Harzsch, S. and Wägele, J.-W. 2004. The brain of the Remipedia (Crustacea) and an alternative hypothesis on their phylogenetic relationships. *Proceedings of the National Academy of Sciences USA* **101**: 3868–3873.
- Farrell, B. D. 1998. “Inordinate Fondness” explained: why are there so many beetles? *Science* **281**: 555–557.
- Fennah, R. G. 1954. The higher classification of the family Issidae (Hemiptera, Fulgoroidea), with description of new species. *Transactions of the Royal Entomological Society* **105**: 455–474.
- 1961. The occurrence of a Cixiinae Fulgoroid (Homoptera) in a Weald Clay. *Annals and Magazine of Natural History* **13**: 161–163.
- 1987. A new genus and species of Cixiidae (Homoptera, Fulgoroidea) from Lower Cretaceous amber. *Journal of Natural History* **21**: 1237–1240.
- Fleck, G., Bechly, G., Martínez-Delclòs, X., Jarzembski, E. A., Coram, R. and Nel, A. 2003. Phylogeny and classification of the Stenophlebiptera. (Insecta, Odonata, Epiproctophora). *Annales de la Société entomologique de France (n.s.)* **39**: 55–93.

- Nel, A., Bechly, G. and Escuillié, F. 2002. The larvae of the Mesozoic family Aeschnidiidae and their phylogenetic implications (Insecta, Odonata, Anisoptera). *Palaeontology* **45**: 165–184.
- Flook, P. K. and Rowell, C. H. F. 1997. The effectiveness of mitochondrial rRNA gene sequences for the reconstruction of the phylogeny of an insect order (Orthoptera). *Molecular Phylogenetics and Evolution* **8**: 177–192.
- and Rowell, C. H. F. 1998. Inferences about orthopteroid phylogeny and molecular evolution from small subunit nuclear ribosomal DNA sequences. *Insect Molecular Biology* **7**: 163–178.
- , Klee, S. and Rowell, C. H. F. 1999. Combined molecular phylogenetic analysis of the Orthoptera (Arthropoda, Insecta) and implications for their higher systematics. *Systematic Biology* **48**: 233–253.
- , Klee, S. and Rowell, C. H. F. 2000. Molecular phylogenetic analysis of the Pneumoroidea (Orthoptera, Caelifera): molecular data resolve morphological character conflicts in the basal Acridomorpha. *Molecular Phylogenetics and Evolution* **15**: 345–354.
- Fontes, L. R. and Vulcano, M. A. 1998. Cupins fósseis do Novo Mundo, pp. 243–295. In Fontes, L. R. and Filho, E. B. (eds), *Cupins – O desafio do conhecimento*. Piracicaba: Fundação de Estudos Agrários Luiz de Queiroz FEALQ.
- and Vulcano, M. A. 2004. Catalog of the Fossil Isoptera of the New World. *Sociobiology* **44**: 345–364.
- Frank, J. H. and Thomas, M. C. 2002. http://creatures.ifas.ufl.edu/misc/beetles/fl_rove_beetles.htm.
- Fraser, F. C. 1957. A reclassification of the order Odonata, based on some new interpretations of the venation of the dragonfly wing. *Handbook Royal Zoological Society of New South Wales* **12**: 1–133.
- Furth, D. G. 1994. Frank Morton Carpenter (1902–1994): academic biography and list of publications. *Psyche* **101**: 127–144.
- Gauld, I. and Bolton, B. 1988. *The Hymenoptera*. British Museum (Natural History). Oxford: Oxford University Press.
- Gebicki, C. and Wegierek, P. 1993. *Oligocixia electrina* gen. et sp. nov. (Homoptera, Auchenorrhyncha, Cixiidae) from Dominican amber. *Annales Naturhistorisches Museum, Wien* **95A**: 121–125.
- Germar, E. F. 1839. Die versteinerten Insekten Solenhofens. *Nova Acta Leopoldina Carolina Akademia* **19**: 187–222.
- Godoy, C. and Webb, M. D. 1994. Recognition of a new subfamily of Cicadellidae from Costa Rica based on a phenetic analysis with similar taxa (Hemiptera Homoptera Auchenorrhyncha). *Tropical Zoology* **7**: 131–144.
- Godunko, R. J. 2004. A new fossil subgenus and species of the genus *Ecdyonurus* Eaton, 1868 from Eocene Baltic amber (Ephemeroptera: Heptageniidae). *Genus* **15**: 323–328.
- and Neumann, C. 2006. Fossil mayfly collections of the Museum für Naturkunde, Humboldt University, Berlin. I. *Electroletus soldani* gen. et sp. nov. (Ephemeroptera: Amelitidae) from the Eocene Baltic Amber.- *Annales Zoologici* **56**: 437–441.
- Gomez-Pallerola, J. E. 1984. Nuevos Paleontíidos del yacimiento Infracretácico de la ‘Pedrera de Meiá’ (Lérida). *Boletín Geológico y Minero España* **95**: 301–309.
- 1986. Nuevos insectos fósiles de las calizas litográficas del Cretacico Inferior del Montsec (Lérida). *Boletín Geológico y Minero España* **97**: 27–46.

- Goodwyn, P. J. P. 2002. A new genus of water measurer from the Lower Cretaceous Crato Formation in Brazil (Insecta: Heteroptera: Gerromorpha: Hydrometridae). *Stuttgarter Beiträge Naturkunde, Serie B* **316**: 1–9.
- Gorochov, A. V. 1992. Fossil stick insects (Phasmatoptera) from the Jurassic and Cretaceous. *Trudy Paleontologicheski Instituta Rossioskyo Akademii, Nauk* 252: 112–116.
- 1994. Permian and Triassic walking sticks (Phasmatoptera) from Eurasia. *Paleontological Journal* **28**: 83–97.
- 1995. The system and evolution of the suborder Ensifera (Orthoptera). *Proceedings of the Zoological Institute, Russian Academy of Sciences* **260**(1): 1–223, **260**(2): 1–212.
- 2000. Phasmomimidae: are they Orthoptera or Phasmatoptera? *Paleontological Journal* **34**: 295–300.
- 2001. The most interesting finds of orthopteroid insects at the end of the 20th century and a new recent genus and species. *Journal of Orthoptera Research* **10**: 353–367.
- and Rasnitsyn, A. P. 2002. Superorder Gryllidea Laicharting, 1781 (= Orthopteroidea Handlirsch, 1903), pp. 293–303. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
- Goulet, H. and Huber, J. T. 1993. *Hymenoptera of the World: an Identification Guide to Families*. Research Branch Agriculture Canada Publication 1894/EL.
- Gratshev, V. G. and Zherikhin, V. V. 1993. New fossil mantids (Insecta, Mantida). *Paleontological Journal* **27**: 148–165.
- and Zherikhin, V. V. 2000. New Early Cretaceous weevil taxa from Spain (Coleoptera, Curculionidae). *Acta Geologica Hispanica* **35**: 37–46.
- Grimaldi, D. 1990. Insects from the Santana Formation, Lower Cretaceous, of Brazil. *Bulletin of the American Museum of Natural History* **195**: 5–191.
- 1991. The Santana Formation insects, pp. 379–406. In Maisey, J. G. (ed), *Santana Fossils: an Illustrated Atlas*. Neptune City, NJ: T.F.H. Publications.
- 1996. *Amber – A window to the past*. AMNH; New York; 216 pp.
- 1999. The co-radiations of pollinating insects and angiosperms in the Cretaceous. *Annals of the Missouri Botanical Garden* **86**: 373–406.
- 2001. Insect evolutionary history from Handlirsch to Hennig, and beyond. *Journal of Paleontology* **75**: 1152–1160.
- 2003. A revision of Cretaceous mantises and their relationships, including new taxa (Insecta: Dictyoptera: Mantodea). *American Museum Novitates* **3412**: 1–47.
- and Maisey, J. G. 1990. Introduction, pp. 5–14. In Grimaldi, D. A. (ed), *Insects from the Santana Formation, Lower Cretaceous, of Brazil*. *Bulletin of the American Museum of Natural History*, **195**.
- and Cumming, J. 1999. Brachyceran Diptera in Cretaceous ambers and Mesozoic diversification of the Eremoneura. *Bulletin of the American Museum of Natural History* **239**: 1–124.
- and Engel, M. S. 2005. *Evolution of Insects*. Cambridge: Cambridge University Press.
- Engel, M. S. and Nascimbene, P. C. 2002. Fossiliferous Cretaceous amber from Myanmar (Burma): its rediscovery, biotic diversity, and paleontological significance. *American Museum Novitates* **3361**: 1–71.
- Guiglia, D. 1955. Appunti intorno al Gen. *Fedtschenkia* Saussure (Hymenoptera: Sapygidae). *Doriania Suppl. Agli* **II** (55): 1–4.
- 1969. The history of the peculiar Genus *Fedtschenkia* Saussure (1880) (Hymenoptera: Fedtschenkiidae). *Israel Journal of Entomology* **4**: 339–342.

- 1972. De la distribution géographique du genre *Fedtschenkia* (Saussure, 1880) (*Hymenoptera: Fedtschenkiidae*). *Proceedings of the XIII Congress of Entomology, Moscow* (1968) **1**: 138–140.
- Gullan, P. J. and Cranston, P. S. 2005. *The Insects – an Outline of Entomology*. Oxford: Blackwell Publishing.
- Gusenleitner, J. 1994. Das Vorkommen der Familie Sapygidae in Österreich. (Insecta: Hymenoptera: Sapygidae). *Annalen des Naturhistorischen Museums in Wien* **96B**: 173–188.
- 1996. Beitrag zur Kenntnis paläarktischer Sapyginae (Hymenoptera, Sapygidae). *Linzer biologische Beiträge* **28**: 23–38.
- 1997. Über Funde von Sapygidae aus dem östlichen Mittelmeergebiet (Sapygidae, Hymenoptera). *Linzer biologische Beiträge* **29**: 105–108.
- Haas, F. and Gorb, S. 2004. Evolution of locomotory attachment pads in the Dermaptera (Insecta). *Arthropod Structure and Development* **33**: 45–66.
- , Gorb, S. and Wootton, R. J. 2000. Elastic joints in dermapteran hind wings: Materials and wing folding. *Arthropod Structure and Development* **29**: 137–146.
- and Kukalová-Peck, J. 2001. Dermaptera hind wing structure and folding: new evidence for familial, ordinal and superordinal relationships within Neoptera (Insecta). *European Journal of Entomology* **98**: 445–509.
- and Klass, K.-D. 2003. The basal phylogenetic relationships in the Dermaptera, pp. 138–142. In Klass, K.-D. (ed), *Proceedings of the 1st Dresden Meeting on Insect Phylogeny: Phylogenetic Relationships within the Insect Orders*, Dresden, 19–21, September 2003. *Entomologische Abhandlungen* **61**.
- Haase, E. 1890. Beiträge zur Kenntnis der fossilen Arachniden. *Zeitschrift der deutschen geologischen Gesellschaft* **42**: 629–657.
- Hamilton, K. G. A. 1971. A remarkable fossil homopteran from Canadian Cretaceous amber representing a new family. *Canadian Entomologist* **103**: 943–946.
- 1972. The insect wing, Part III. Venation of the orders. *Journal of the Kansas Entomological Society* **45**: 145–162.
- 1983. Classification, morphology and phylogeny of the family Cicadellidae (Rhynchota: Homoptera), pp. 15–37. In Knight, W. J. (ed), *International Workshop on Leafhoppers and Planthoppers of Economic Importance*. London: Commonwealth Institute of Entomology.
- 1990. Homoptera, pp. 82–122. In Grimaldi, D. (ed), Insects from the Santana Formation, Lower Cretaceous, of Brazil. *Bulletin of the American Museum of Natural History* **195**.
- 1992. Lower Cretaceous Homoptera from the Koonwarra Fossil Bed in Australia with a New Superfamily and Synopsis of Mesozoic Homoptera. *Annals of the Entomological Society of America* **85**: 423–430.
- 1996. Cretaceous Homoptera from Brazil: Implications for Classification, pp. 89–110. In Schaefer, C. W. (ed), *Studies on Hemipteran Phylogeny*. Langham, MD: Thomas Say Publications in Entomology. Entomological Society of America.
- 1999. The ground-dwelling leafhoppers Myerslopiidae, new family and Sagmatiini, new tribe (Homoptera: Membracoidea). *Invertebrate Taxonomy* **13**: 207–235.
- Handlirsch, A. 1906–1908. *Die fossilen Insekten und die Phylogenie der rezenten Formen. Ein Handbuch für Paläontologen und Zoologen*. Leipzig: Engelmann.
- 1925. Palaeontology, pp. 117–306. In Schröder, C. (ed), *Handbuch der Entomologie* **3**. Jena: G. Fischer.
- 1939. Neue Untersuchungen über die fossilen Insekten mit Ergänzungen und Nachträgen sowie Ausblicken auf phylogenetische, palaeogeographische und

- allgemein biologische Probleme, II Theil. *Annalen des Naturhistorischen Museums in Wien* **48**: 1–240.
- Hasegawa, E. and Kasuya, E. 2006. Phylogenetic analysis of the insect order Odonata using 28S and 16S rDNA sequences: a comparison between data sets with different evolutionary rates. *Entomological Science* **9**: 55–66.
- Hasiotis, S. T. and Dubiel, R. F. 1995. Termite (Insecta: Isoptera) nest ichnofossils from the Triassic Chinle Formation, Petrified Forest National Park, Arizona. *Ichnos* **4**: 119–130.
- , Peterson, F., Demko, T. M. and Turner, C. 1997. Giant termite (Insecta: Isoptera) nests from the base of the Upper Jurassic Morrison Formation, northwestern New Mexico. *Geological Society of America Abstracts with Program* **28**: A461.
- Heads, S. W. 2008. The first fossil Proscopiidae (Insecta, Orthoptera, Eumastacidae) with comments on the historical biogeography and evolution of the family. *Palaeontology* (in press).
- Martill, D. M. and Loveridge, R. F. 2005. An exceptionally preserved antlion (Insecta, Neuroptera) with colour pattern preservation from the Cretaceous of Brazil. *Palaeontology* **48**: 1409–1417.
- Hennig, W. 1953. Kritische Bemerkungen zum phylogenetischen System der Insekten. *Beiträge zur Entomologie* **3**: 1–85.
- 1954. Flügelgeäder und System der Dipteren. *Beiträge zur Entomologie* **4**: 245–388.
- 1969. *Die Stammesgeschichte der Insekten*. Frankfurt: W. Kramer.
- 1973. Diptera (Zweiflügler). In *Handbuch der Zoologie*, IV. Band: Arthropoda – 2. Hälfte: Insecta – 2. Teil: Spezielles, 31. Berlin: Walter de Gruyter.
- 1981. *Insect Phylogeny*. Chichester: Wiley.
- Herczek, A. and Popov, Yu. A. 2001. Redescription of the oldest plant bugs from the Upper Jurassic of the Southern Kazakhstan (Heteroptera: Cimicomorpha, Miridae). *Annals of the Upper Silesian Museum in Bytom, Entomology* **10–11**: 121–128.
- Herman, L. H. 2001. Catalog of the Staphylinidae (Insecta: Coleoptera). 1758 to the end of the second millennium. I. introduction, history, biographical Sketches, and omaliine group. *Bulletin of the American Museum of Natural History* **265**: 1–650.
- Hoch, H., Deckert, J. and Wessel, A. 2006. Vibrational signalling in a Gondwanan relict insect (Hemiptera: Coleorrhyncha: Peloridiidae). *Biology Letters* **2**: 222–224.
- Hölldobler, B. and Wilson, E. O. 1990. *The Ants*. Cambridge: Belknap Press of Harvard University Press.
- Holzinger, W. E., Kammerlander, I., Bourgoin, Th., Chan, K. L. and Campbell, B. C. 2001. Towards a phylogeny of the Cixiidae (Fulgoromorpha) and its major subgroups: preliminary results. *2nd European Hemiptera Congress*, 20–24 June 2001, Fiesa, Slovenia: 19.
- , Emeljanov, A. F. and Kammerlander, I. 2002. The family Cixiidae – a review, pp. 113–138. In Holzinger, W. (ed), *Zikaden – Leafhoppers, Planthoppers and Cicadas (Insecta: Hemiptera: Auchenorrhyncha)*. Denisia, 04, Zugleich Kataloge Neue Folge Nr. 176, Linz: OÖ. Landesmuseums.
- Hong, Y. 1982. *Mesozoic Fossil Insects of Jiuquan Basin, Gansu Province*. Beijing: Geological Publishing House.
- Hörnschemeyer, T. 1994. Ein fossiler Tenebrionidae *Ceropria? messelense* n. sp. (Coleoptera: Tenebrionidae: Diaperinae) aus dem Mitteleozän der Grube Messel bei Darmstadt. *Courier Forschungsinstitut Senckenberg* **170**: 75–83.
- Hovmöller, R. 2006. *Molecular phylogenetics and taxonomic issues in dragonfly systematics (Insecta: Odonata)*. PhD thesis, Stockholm University.

- Hubbard, M. D. and Kukalová-Peck, J. 1980. Permian mayfly nymphs: new taxa and systematic characters, pp. 19–31. In Flannagan, F. F. and Marshall, K. E. (eds), *Advances in Ephemeroptera Biology*. New York: Plenum Press.
- and Savage, H. M. 1981. The fossil Leptophlebiidae (Ephemeroptera): A systematic and phylogenetic review. *Journal of Paleontology* **55**: 810–813.
- Hughes, L. and Westoby, M. 1992. Capitula on stick insect eggs and elaiosomes on seeds: convergent adaptations for burial by ants. *Functional Ecology* **6**: 642–648.
- Illies, J. 1968. Ephemeroptera (Eintagsfliegen). *Handbuch der Zoologie IV (Arthropoda) 2 (Insecta)* **5**: 1–63.
- Ivanov, V. D. and Sukatsheva, I. D. 2002. Order Trichoptera Kirby, 1813. The Caddisflies, pp. 199–220. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
- James, M. T. 1981. Stratiomyidae, pp. 497–511. In McAlpine, J. F. and Woods, D. M. (eds), *Manual of Nearctic Diptera*. Ottawa: Biosystematics Research Institute.
- Jarvis, K. J., Haas, F. and Whiting, M. 2005. A phylogeny of earwigs (Insecta: Dermaptera) based on molecular and morphological evidence: reconsidering the classification of Dermaptera. *Systematic Entomology* **30**: 442–453.
- Jarzembski, E. A. 1991. New insects from the Weald Clay of the Weald. *Proceedings of the Geologists' Association* **9**: 71–93.
- , Martínez-Delclòs, X., Bechly, G., Nel, A., Coram, R. and Escuillié, F. 1998. The Mesozoic non-calopterygoid Zygoptera: description of new genera and species from the Lower Cretaceous of England and Brazil and their phylogenetic significance (Odonata, Zygoptera, Coenagrionoidea, Hemiphlebioidea, Lestoidea). *Cretaceous Research* **19**: 403–444.
- Jell, P. A. and Duncan, P. M. 1986. Invertebrates, mainly insects, from the freshwater Lower Cretaceous, Koonwarra Fossil Bed (Korumburra Group), South Gippsland, Victoria, pp. 111–205. In *Plants and Invertebrates from the Lower Cretaceous Koonwarra Fossil Bed, South Gippsland, Victoria. Memoirs of the Association of Australasian Palaeontologists* **3**.
- Kaszab, Z. and Schawaller, W. 1984. Eine neue Schwarzkäfer-Gattung und -Art aus Dominikanischem Bernstein (Coleoptera, Tenebrionidae). *Stuttgarter Beiträge zur Naturkunde, Serie B* **109**: 1–6.
- Key, K. H. L. 1991. Phasmatodea, pp. 394–403. In Naumann, D. (ed), *The Insects of Australia: a Textbook for Students and Research Workers*, vol. 1, 2nd edn. Carlton: Melbourne University Press.
- Kimsey, L. S. 1991. Relationships among the tephid wasp subfamilies (Hymenoptera). *Systematic Entomology* **16**: 427–438.
- Kirejtshuk, A. G. and Ponomarenko, A. G. 1990. Fossil beetles of the Peltidae and Nitidulidae families (Coleoptera). *Paleontological Journal* **24**: 79–90.
- Klass, K. D. 1997a. The external male genitalia and the phylogeny of Blattaria and Mantodea. *Bonner Zoologische Monographien* **42**: 1–341.
- 1997b. The ovipositor of Dictyoptera (Insecta): Homology and ground-plan of the main elements. *Zoologischer Anzeiger* **236**: 69–101.
- 2001. Morphological evidence on Blattaria phylogeny: “Phylogenetic histories and stories” (Insecta, Dictyoptera). *Deutsche Entomologische Zeitschrift* **48**: 223–265.
- Kluge, N. J. 1986. Recent mayfly species (Ephemeroptera, Heptageniidae) in Baltic amber. *Paleontological Journal* **2**: 111–112.
- 1996. A new suborder of Thysanura for the Carboniferous insect originally described as larva of *Bojophlebia*, with comments on characters of the orders Thysanura and Ephemeroptera. *Zoosystematica Rossica* **4**: 71–75.

- 1998. Phylogeny and higher classification of Ephemeroptera. *Zoosystematica Rossica* **7**: 255–269.
- 2004. *The Phylogenetic System of Ephemeroptera*, Dordrecht: Kluwer Academic Publishers.
- , Studemann, D., Landolt, P. and Gonser, T. 1995. A reclassification of Siphlonuroidea (Ephemeroptera). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* **68**: 103–32.
- , Godunko, R. J. and Krzeminski, W. 2006. A new mayfly family (Insecta: Ephemeroptera) from Eocene Baltic amber. *Annales Zoologici* **56**: 181–185.
- Koch, M. 1997. Monophyly and phylogenetic position of the Diplura (Hexapoda). *Pedobiologia* **41**: 9–12.
- 2003. Towards a phylogenetic system of the Zygentoma. *Entomologische Abhandlungen* **61**: 122–124.
- Königsmann, E. 1976. Das phylogenetische System der Hymenoptera, Sapygidae. *Deutsche Entomologische Zeitschrift N.F.* **25(IV–V)**: 406–409.
- Kozlov, M. V., Ivanov, V. D. and Rasnitsyn, A. P. 2002. Order Lepidoptera Linné, 1758. The butterflies and moths, pp. 220–227. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
- Krell, F.-T. 2000. The fossil record of Mesozoic and Tertiary Scarabaeoidea (Coleoptera: Polyphaga). *Invertebrate Taxonomy* **2000**: 871–905.
- Krishna, K. 1990. Isoptera, pp. 76–81. In Grimaldi, D. A. (ed), *Insects from the Santana Formation, Lower Cretaceous, of Brazil. Bulletin of the American Museum of Natural History* **195**.
- Kristensen, N. P. 1991. Phylogeny of extant hexapods, pp. 125–140. In Naumann, I. D. (ed), *The Insects of Australia: a Textbook for Students and Research Workers*, vol. 1, 2nd edn. Melbourne: Melbourne University Press.
- 1995. Forty years' insect phylogenetic systematics: Hennig's "Kritische Bemerkungen" and subsequent developments. *Zoologische Beiträge (NF)* **36**: 83–124.
- 1998. The groundplan and basal diversification of the hexapods, pp. 281–293. In Fortey, R. A. and Thomas, R. H. (eds), *Arthropod Relationships*. London: Chapman and Hall.
- Krzeminski, W. and Evenhuis, N. L. 2000. Review of Diptera paleontological records, pp. 535–564. In Papp, L. and Darvas, B. (eds), *Contributions to a Manual of Palaearctic Diptera*, vol. 1 L. Budapest: Science Herald.
- Kukalová-Peck, J. 1978. Origin and evolution of insect wings and their relation to metamorphosis, as documented by the fossil record. *Journal of Morphology* **156**: 53–125.
- 1983. Origin of the insect wing and wing articulation from the arthropodan leg. *Canadian Journal of Zoology* **61**: 1618–1669.
- 1985. Ephemeroptery wing venation based upon new gigantic Carboniferous mayflies and basic morphology, phylogeny, and metamorphosis of pterygote insects (Insecta, Ephemeroptera). *Canadian Journal of Zoology* **63**: 933–955.
- 1987. New Carboniferous Diplura, Monura, and Thysanura, the hexapod ground plan, and the role of thoracic side lobes in the origin of wings (Insecta). *Canadian Journal of Zoology* **65**: 2327–2345.
- 1991. Fossil history and the evolution of hexapod structures, pp. 141–179. In Naumann, I. D. (ed), *The Insects of Australia: a Textbook for Students and Research Workers*, vol. 1. Melbourne: Melbourne University Press.
- Kuschel, G. 2003. Nemonychidae, Belidae, Brentidae (Insecta: Coleoptera: Curculionoidea). *Fauna of New Zealand* **45**: 1–100. Manaaki Whenua Press.

- Kwok, H. K. and Corlett, R. T. 2002. Seasonality of forest invertebrates in Hong Kong, South China. *Journal of Tropical Ecology* **18**: 637–644.
- Ladiges, P. Y., Nelson, G. and Grimes, J. 1997. Subtree analysis, *Nothofagus* and Pacific biogeography. *Cladistics* **13**: 125–129.
- Lameere, A. 1917. Étude sur l'évolution des Ephémères. *Bulletin de Societe Zoologique de France* **42**: 41–81.
- Lauterbach, K. E. 1973. Schlüsselereignisse in der Evolution der Stammgruppe der Euarthropoda. *Zoologische Beiträge, Neue Folge* **19**: 251–299.
- Lawrence, J. F. 1999. The Australian Ommatidae (Coleoptera: Archostemata): new species, larva and discussion of relationships. *Invertebrate Taxonomy* **13**: 369–390.
- and Newton, Jr. A. F. 1995. Families and subfamilies of Coleoptera (with selected genera, notes, references and data on family-group names), pp. 779–1006. In Pakaluk, J. and Slipinski, S. A. (eds), *Biology, Phylogeny, and Classification of Coleoptera*, vol. 2. Warsaw: Muzeum I Instytut Zoologii PAN.
- Lin, Q.-B., and Huang, D. Y. 2001. Description of *Caenoephemera shangyuensis*, gen. nov., sp. nov. (Ephemeroptera), from the Yixian Formation. *Canadian Entomologist* **133**: 747–754.
- Nel, A. and Huang, D.-Y. 2002. Phylogenetic analysis of the Mesozoic dragonfly family Liupanshaniidae (Insecta: Aeshnoptera: Odonata). *Cretaceous Research* **23**: 439–444.
- Lo, N., Bandi, C., Watanabe, H., Nalepa, Ch. and Beninati, T. 2003. Evidence for coeladogenesis between diverse dictyopteran lineages and their intracellular endosymbionts. *Molecular Biology and Evolution* **20**: 907–913.
- Lohmann, H. 1996. Das Phylogenetische System der Anisoptera (Odonata). *Entomologische Zeitschrift* **106(6)**: 209–252; **106(7)**: 253–296; **106(9)**: 360–367.
- Lomhold, O. 1982. On the origin of the bees (Hymenoptera: Apidae, Sphecidae). *Entomologica Scandinavica* **13**: 185–190.
- Lubkin, S. H. 2003. *Paracupes svitkoi* (Coleoptera: Cupedidae), a new species from the Cretaceous of New Jersey. *Acta Zoologica Cracoviensia* **46** (suppl. Fossil Insects): 189–194.
- and Engel, M. S. 2005. *Permocoleus*, New Genus, the First Permian Beetle (Coleoptera) from North America. *Annals of the Entomological Society of America* **98**: 73–76.
- Maddison, D. R. 1995. Coptoclavidae. <http://tolweb.org/tree/tree?group=Coptoclavidae>.
- Maisey, J. G. (ed.). 1991. *Santana Fossils: an Illustrated Atlas*. Neptune City, NJ: T.F.H. Publications.
- Makarkin, V. N. and Menon, F. 2007. First record of fossil ‘rapismatid-like’ Ithionidae (Insecta, Neuroptera) from the Lower Cretaceous Crato Formation of Brazil. *Cretaceous Research* (in press).
- Malyshev, S. I. 1968. *Genesis of the Hymenoptera and the Phases of their Evolution*. London Methuen & Co.
- Marden, J. H. and Thomas, M. A. 2005. Rowing locomotion by a stonefly that possesses ancestral pterygote condition of co-occurring wings and abdominal gills. *Biological Journal of the Linnean Society* **79**: 341–349.
- Marshall, J. A and Haes, E. C. M. 1988. *Grasshoppers and allied insects of Great Britain and Ireland*. Colchester: Harley Books.
- Martill, D. M. (ed.) 1993. *Fossils of the Santana and Crato Formations, Brazil*. The Palaeontological Association Field Guides to Fossils, 5. London: The Palaeontological Association.

- 1994. Fake fossils from Brazil. *Geology Today* May/June **1994**: 111–115.
- and Frey, E. 1995. Colour patterning preserved in Lower Cretaceous birds and insects: the Crato-Formation of N.E. Brazil. *Neues Jahrbuch für Geologie und Paläontologie Monatshefte* **2**: 118–128.
- and Nel, A. 1996. A new dragonfly from the Crato Formation (Lower Cretaceous, Aptian) of N. E. Brazil. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* **5**: 279–292.
- and Davis, P. G. 1998. Did dinosaurs come up to scratch? *Nature* **396**: 528–589.
- Martínez, S. 1982. Catalogo sistemático de los insectos fosiles de America del Sur. *Revista de la Facultad de Humanidades y Ciencias, Serie Ciencias de la Tierra* **1**: 29–83.
- Martínez-Delclós, X. 1989. *Chresmoda aquatica* n. sp. insecto Chresmodidae del Cretácico inferior de la Sierra del Montsech (Lleida, España). *Revista Española de Paleontología* **4**: 67–74.
- , Nel, A., Azar, D., Bechly, G., Dunlop, J. A., Engel, M. S. and Heads, S. 2007. The enigmatic, Mesozoic family Chresmodidae (Polyneoptera: Archaeorthoptera): New palaeobiological and phylogenetic data, with the description of a new species from Brazil – New palaeobiological and phylogenetical data. *Neues Jahrbuch für Geologie und Paläontologie – Abhandlungen* (in press).
- Martins-Neto, R. G. 1987a. Um novo gênero de Orthoptera (Insecta, Grylloidea) da Formação Santana, Bacia do Araripe (Cretáceo Inferior) nordeste do Brasil. *Anais 10º Congresso Brasileiro de Paleontologia*, Rio de Janeiro: 599–609.
- 1987b. A paleoentomofauna Brasileira: estagio actual do concimento. *Anais 10º Congresso Brasileiro Paleontologia*, Rio de Janeiro v. **II**: 567–591.
- 1987c. Descrição de três novos géneros e três novas espécies de Orthoptera (Insecta, Acridoidea) da Formação Santana, Bacia do Araripe (Cretáceo Inferior) nordeste do Brasil, representando três familiias, sendo que duas novas: Archaeopneumoridae nov. fam. e Bouretidae nov. fam. *Anais da Academia Brasileira de Ciências* **59**: 444.
- 1988a. A new fossil insect (Homoptera, Cixiidae) from the Santana Formation (Lower Cretaceous), Araripe Basin, Northeast Brasil. *Interciencia* **13**: 313–316.
- 1988b. Neuropteros (Insecta: Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, nordeste do Brasil. I. Família Chrysopidae. *Anais da Academia Brasileira de Ciências* **60**: 189–201.
- 1989a. A new fossil insect (Homoptera, Cixiidae) from the Santana Formation (Lower Cretaceous) Araripe Basin, Northeast Brazil. *Acta Geologica Leopoldensia* **26**: 7–14.
- 1989b. Primeiro registro de Phasmatodea (Insecta: Orthopteromorpha) na Formação Santana, Bacia do Araripe (Cretáceo Inferior), nordeste do Brasil. *Acta Geologica Leopoldensia (Estudos Tecnológicos)* **12**: 91–104.
- 1990a. Primeiro registro de Dermaptera (Insecta, Orthopteromorpha) na Formação Santana (Cretáceo inferior), Bacia do Araripe, nordeste do Brasil. *Revista Brasileira de Entomologia* **34**: 775–784.
- 1990b. Neurópteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. VI. Ensaio filogenético das espécie do genera *Blittersdorffia* Martins-Neto and Vulcano, com descrição de nova espécie. *Acta Geologica Leopoldensia* **13**: 3–12.
- 1990c. The family Locustopsidae (Insecta, Caelifera) in the Santana Formation (Lower Cretaceous, northeast Brazil). I. – Description of two new species of the genus *Locustopsis* Handlirsch and three new species of the genus *Zessinia* n. gen. *Atlas do I Simpósio sobre a Bacia do Araripe e Bacias Interiores do n Nordeste Crato*, 14 a 16 Junho de 1990: 277–91.

- 1990d. Um novo gênero e duas nova espécies de Tridactylidae (Insecta, Orthopteroidea) na Formação Santana (Cretáceo Inferior do nordeste do Brasil). *Annales Academia Brasil Ciencias* **62**: 51–59.
- 1991a. Sistemática dos Ensifera (Insecta, Orthopteroida) da Formação Santana, Cretáceo Inferior do nordeste do Brasil. *Estudio Technológico, Acta Geologica Leopoldensia* **32**: 3–162.
- 1991b. *Cratogryllus ciguaeli*, nova espécie de Ensifera (Insecta, Grylloidea) da Formação Santana (Cretáceo Inferior) Bacia do Araripe, nordeste do Brasil. *Acta Geologica Leopoldensia* **33**: 153–156.
- 1991c. Evidências de especiação alocrônica na fauna de Ensifera (Insecta, Orthopteroida) da Formação Santana, Cretáceo do nordeste do Brasil. *Revista de Geologia* **4**: 61–80.
- 1991d. A Paleoentomofauna do Nordeste Brasileiro: Estado da Arte. *Atas XIV Simposio de Geologia do Nordeste, Recife* **12**: 59–62.
- 1992a. Neurópteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. V. Aspectos filogenéticos, paleoecológicos, paleobiogeográficos e descrição de novos taxa. *Anais da Academia Brasileira de Ciências* **64**: 117–148.
- 1992b. Neurópteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. VII. Palaeoleontinae, nova subfamília de Myrmeleontidae e descrição de novos táxons. *Revista Brasileira de Entomologia* **36**: 803–815.
- 1992c. Nova ocorrência, variabilidade morfológica e relações filogenéticas do gênero *Cratoelcana* Martins-Neto, 1991 (Insecta, Ensifera, Elcanidae), da Formação Santana, Bacia do Araripe, Brasil. *Revista Brasil Entomologia* **36**: 817–830.
- 1992d. A paleoentomofauna das bacias mesozoicas brasileiras e seu potencial para possíveis correlações bioestratigráficas entre depósitos não-marinhos. In I. Encontro sobre a sedimentação continental das bacias mesozoicas brasileiras. *Acta Geologica Leopoldensia* **36**: 125–126.
- 1994. Neurópteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. IX. Primeiros resultados da composição da fauna e descrição de novos táxons. *Acta Geologica Leopoldensia* **17**: 269–288.
- 1995a. Araripelocustidae, fam. n. uma nova família de gafanhotos (Insecta, Caelifera) da Formação Santana, Cretáceo Inferior do nordeste do Brasil. *Revista Brasil Entomologia* **39**: 311–319.
- 1995b. Complementos ao estudo sobre os Ensifera (Insecta, Orthopteroida) da Formação Santana, Cretáceo Inferior do nordeste do Brasil. *Revista Brasil Entomologia* **39**: 321–345.
- 1996a. New mayflies (Insecta, Ephemeroptera) from the Santana Formation (Lower Cretaceous), Araripe Basin, Northeastern Brazil. *Revista Española de Paleontología* **11**: 177–192.
- 1996b. Interação inseto / flora e considerações sobre a origem das angiospermas. *Revista UnG, Série Geociências* **1**: 31–36.
- 1997a. Neurópteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. X. Descrição de novos táxons (Chrysopidae, Babinskaiidae, Myrmeleontidae, Ascalaphidae e Psychopsidae). *Revista Universidade de Guarulhos, Série Ciências Exatas e Technológicas* **2**: 68–83.
- 1997b. *Cratotetraspinus* nov. nom., novo nome para *Tetraspinus* Martins-Neto, 1995 (nom. preoc.). *Revista Universidade Guarulhos, Geociências* **2**: 105.

- 1998a. A new genus of the family Locustopsidae (Insecta, Caelifera) in the Santana Formation (Lower Cretaceous, northeast Brazil). *Revista Española de Paleontología* **13**: 133–138.
- 1998b. A new subfamily of Ensifera (Insecta, Grylloidea) from the Santana Formation (Lower Cretaceous), Araripe Basin, NE Brazil. *Proceedings of the First International Palaeoentomological Conference*, Moscow 1998: 91–97.
- 1998c. *Conan barbarica* n. gen. et n. sp. (Insecta, Coleoptera, Coptoclavidae) – uma gigantesca larva da Formação Santana, Cretáceo Inferior, Bacia do Araripe, Brasil. *Geociências São Paulo* **17**: 109–114.
- 1998d. Neurópteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. XI. Descrição de novos táxons de Myrmeleontidae (Palaeoleontinae e Pseudonymphinae). *Revista Universidade de Guarulhos, Série Ciências Biológicas e da Saúde* **3**: 38–42.
- 1998e. The paleoentomo fauna from Tremembé Formation (Taubaté Basin) Oligocene of São Paulo State: new Hemiptera, Auchenorrhyncha, Hymenoptera, Coleoptera, Lepidoptera and Diptera (Parte II) (Insecta). *Revista Universidade Guaralhos, Serie Geociências* **3**: 58–70.
- 1998f. Novos registro de Palaeointinideos (Insecta: Hemiptera) na formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. *Acta Geologica Leopoldensia* **21**: 69–74.
- 1999a. Estado actual del conocimiento de la paleoentomofauna brasileña. *Revista de la Sociedad Entomologica Argentina* **58**: 71–85.
- 1999b. New genus and new species of Lepidoptera (Insecta, Eolepidopterigidae) from Santana Formation (Lower Cretaceous, northeast Brazil). *5º Simpósio sobre o Cretáceo do Brasil*, Rio Claro. Boletim: 531–535.
- 2000. Remarks on the neuropterofauna (Insecta, Neuroptera) from the Brazilian Cretaceous, with keys for the identification of the known taxa. *Acta Geologica Hispanica* **35**: 97–118.
- 2001a. Review of some Insecta from Mesozoic and Cenozoic Brazilian deposits with descriptions of new taxa. *Acta Geologica Leopoldensia* **24**: 115–124.
- 2001b. Primeiro registro de Trichoptera (Insecta) na Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil, com descrição de sete novos táxons. *Simpósio sobre a Bacia do Araripe e bacias interiores do Nordeste*, 1 e 2, Crato, 1990/1997. Boletim: 212–226.
- 2002a. The Santana Formation Paleoentomofauna reviewed. Part I. Neuropteroida (Neuroptera and Raphidioptera): systematic and phylogeny, with descriptions of new taxa. *Acta Geologica Leopoldensia* **25**: 35–66.
- 2002b. *Insetos fósseis como bioindicadores em depósitos sedimentares: um estudo de caso para o mesozóico sul-americano*. São Leopoldo: CPGEO, UNISINOS.
- 2003a. Systematics of the Caelifera (Insecta, Orthopteroidea) from the Santana Formation, Araripe Basin (Lower Cretaceous, northeast Brazil), with a description of new genera and species. *Acta Zoologica Cracoviensis* **46** (suppl.): 205–228.
- 2003b. The fossil Tabanids (Diptera Tabanidae): When they began to appreciate warm blood and when they began transmit diseases? *Memorias do Instituto Oswaldo Cruz* **98** (suppl. I): 29–34.
- 2004. New Neuroptera (Nymphidae and Araripeneuridae) from Santana Formation (Lower Cretaceous, Araripe Basin, Northeast Brasil). *Acta Geologica Leopoldensia* **27**.

- 2005a. The Santana Formation palaeoentomofauna reviewed: Polyneoptera (Isoptera, Blattoptera, Dermaptera, Orthoptera) with description of six new species, one new genus and one new combination. *3rd International Congress of Palaeoentomology, Abstracts*, 7–11 February 2005, Pretoria, South Africa. **1**: 24.
- 2005b. Estágio atual da paleoartropologia brasileira: Hexápodes, miriápodes, crustáceos (Isopoda, Decapoda, Eucrustacea e Copepoda) e quelicerados. *Arquivos do Museu Nacional, Rio de Janeiro* **63**: 471–494.
- and Vulcano, M. A. 1989a. Primeiro registro de Raphidioptera (Neuropteroidea) na Formação Santana (Cretáceo Inferior), Bacia do Araripe, nordeste do Brasil. *Revista Brasileira de Entomologia* **34**: 241–249.
- and Vulcano, M. A. 1989b. Amphiesmenoptera, (Trichoptera + Lepidoptera) na Formação Santana (Cretáceo Inferior) Bacia do Araripe, Nordeste do Brasil. I – Lepidoptera (Insecta). *Anais da Academia Brasileira de Ciências* **61**: 459–465.
- and Vulcano, M. A. 1989c Neuropteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. I. Família Chrysopidae. *Anais Academia Brasileira de Ciências* **60**: 189–201.
- and Caldas, E. B. 1990. Efêmeras escavadoras (Insecta, Ephemeroptera, Ephemeroidea) na Formação Santana (Cretáceo Inferior), Bacia do Araripe Nordeste do Brasil: Descrição de três novos gêneros e três novas espécies (ninfas). *Atas do I Simpósio sobre a Bacia do Araripe e Bacias Inteiores do Nordeste Crato*, 1990: 265–275.
- and Vulcano, M. A. 1990a. Neuropteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, nordeste do Brasil. III – Superfamília Mantispoidea. *Revista Brasileira de Entomologia* **34**: 619–625.
- and Vulcano, M. A. 1990b. Neuropteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, nordeste do Brasil. VI – Ensaio filogenético das espécies do gênero *Blittersdorfia*, com descrição de nova espécie. *Acta Geologica Leopoldensia* **31**: 3–12.
- and Valarelli, J. V. 1991. Primeiro registro de Eumastacoidea (Insecta, Caelifera) da Formação Santana, Cretáceo Inferior do nordeste do Brasil. *Anais da Academia Brasileira de Ciências* **63**: 91–92.
- and Nel, A. 1992. Un nouveau fossile de Raphidoptère de la formation Santana, Crétacé inférieur du Brésil (Neuropteroidea, Raphidoptera). *Bulletin de la Société Entomologique de France* **97**: 425–428.
- and Santos, J. C. K. 1994. Um novo gênero e uma nova espécie de Mutuca (Insecta, Diptera, Tabanidae) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. *Acta Geologica Leopoldensia* **39**: 289–297.
- and Vulcano, M. A. 1997. Neuropteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, nordeste do Brasil. VIII – Descrição de novos taxa de Myrmeleontidae, Ascalaphidae e Nemopteridae. *Revista Universidade Guarulhos, Série Ciências Biológicas* **2**: 64–81.
- and Petrulevičius, J. F. 1998. Mesozoic and Cenozoic Orthopteroida from South America. *1st Paleoentomological Conference, Abstracts*, 30 August–4 September 1998, Moscow: 24.
- and Szwedo, J. 2007. Taxonomic changes in fossil Cixiidae (Insecta: Hemiptera: Fulgoromorpha) 1 – *Cretofennahia* nom. nov. pro *Fennahia* Martins-Neto. *Alevisia* **1**: 117.
- , Popov, Yu. A., and Zamboni, J. A. 1999. First South Hemisphere Cretaceous record of Coreoidea (Insecta, Heteroptera) from Santana Formation (Lower Cretaceous, Northeast Brazil), representing a new genus et species. *Bulletim do 5^o Simposio Sobre o Cretáceo do Brasil*: 525–530.

- , Ribeiro-Júnior, C. and Prezoto, F. 2006. New fossils (Isoptera: Hodotermitidae), from the Santana Formation (Lower Cretaceous, Araripe Basin, Northeast Brazil), with descriptions of new taxa including a new subfamily. *Sociobiology* **47**: 125–134.
- , Melo, A. C. and Prezoto, F. 2007. A new species of wasp (Symphyta, Sepulcidae) from the Santana Formation (Lower Cretaceous, Northeast Brazil). *Journal of the Entomological Research Society* **9(1)**: 1–6.
- Martynov, A. V. 1925. Über zwei Grundtypen der Flügel bei den Insecten und ihre Evolution. *Zeitschrift zur Morphologie und Ökologie der Tiere* **4**: 465–501.
- Martynova, O. M. 1962 (reprinted 1991). Order Phasmatodea, pp. 213–215. In Rohdendorf, B. B. (ed), *Fundamentals of Paleontology*, vol. 9, *Arthropoda, Tracheata, Chelicerata*. Washington DC: Smithsonian Institution Libraries and The National Science Foundation.
- Matzke, D. and Klass, K.-D. 2005. Reproductive biology and nymphal development in the basal earwig *Tagalina papua* (Insecta: Dermaptera: Pygidicranidae), with a comparison of brood care in Dermaptera and Embioptera. *Entomologische Abhandlungen* **62**: 99–116.
- Mazzarolo, L. A. and Amorim, D. S. 2000. *Cratomyia macrorrhyncha*, a Lower Cretaceous brachyceran fossil from the Santana Formation, Brazil, representing a new species, genus and family of the Stratiomyomorpha (Diptera). *Insect Systematics & Evolution* **31**: 91–102.
- Mazzoni, A. F. and Hünicken, M. A. 1984. Ontogenia de los notonectidos (Insecta, Heteroptera) del Cretácico inferior de San Luis, Argentina. *Memoria III Congreso Latinoamericano de Paleontología*: 388–395.
- and Hünicken, M. A. 1987. Corixidae (Insecta, Heteroptera) del Cretácico Inferior de la Sierra del Gigante, San Luis, Argentina. *Actas IV Congresso Latinoamericano Palaeontologia, Bolivia* **2**: 731–738.
- McCafferty, W. P. 1990. Ephemeroptera, pp. 20–50. In Grimaldi, D. A. (ed), Insects from the Santana Formation, Lower Cretaceous, of Brazil. *Bulletin of the American Museum of Natural History* **195**.
- 1991. Toward a phylogenetic classification of the Ephemeroptera (Insecta): a commentary on systematics. *Annals of the Entomological Society of America* **84**: 343–360.
- 1997. Ephemeroptera, pp. 89–117. In Poole, R. W. and Gentilli, P. (eds), *Nomina Insecta Nearctica, a Checklist of the Insects of North America*, vol. 4. *Non-holometabolous Orders*. Rockville, MD: Entomological Informational Service.
- McCafferty, W. P. 1998. Ephemeroptera and the great American interchange. *Journal of the North American Benthological Society* **17**: 1–20.
- 2004. Higher classification of the burrowing mayflies (Ephemeroptera: Shappodonta). *Entomological News* **115**: 84–92.
- and Edmunds, Jr. G. F. 1979. The higher classification of the Ephemeroptera and its evolutionary basis. *Annals of the Entomological Society of America* **72**: 5–12.
- McAlpine, J. F. 1981. Morphology and terminology – adults, pp. 9–63. In McAlpine, J. F. and Woods, D. M. (eds), *Manual of Nearctic Diptera*, vol. 1. Ottawa: Biosystematics Research Institute.
- Melo, G. A. R. 1999. Phylogenetic relationships and classification of the major lineages of Apoidea (Hymenoptera), with emphasis on the crabronid wasps. *Scientific Papers of the Natural History Museum, University of Kansas* **14**: 1–55.
- Mendes, L. F. 1997a. A first contribution to the study of the Dominican amber Zygentoma (Insecta), family Ateluridae. *Pedobiologia* **41**: 40–43.
- 1998a. Second contribution to the study of the Dominican amber Zygentoma (Insecta): Family Lepismatidae. *Canadian Entomologist* **130**: 899–904.

- Mendes, M. 1991a. A utilização de caracteres morfológicos alternativos na Sistemática dos Blattoptera (Insecta) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. *XII Congresso Brasileiro de Paleontologia, 1991, São Paulo, Boletim de Resumos 1*: 54.
- 1991b. Raphidiomimidae (Insecta, Blattoptera) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. *XII Congresso Brasileiro de Paleontologia, 1991, São Paulo, Boletim de Resumos 1*: 55.
- 1993. Um novo gênero da Família Blattidae (Insecta, Blattoptera) na Formação Santana, Cretáceo do Nordeste do Brasil. *XIII Congresso Brasileiro de Paleontologia, 1993, São Leopoldo, Boletim de Resumos 1*: 115.
- 1995. Três Novas Espécies do Gênero Mesoblattina (Insecta, Blattoptera) na Formação Santana, Cretáceo do Nordeste do Brasil. *XIV Congresso Brasileiro de Paleontologia, 1995, Uberaba, Boletim de Resumos 1*: 98.
- 1997b. Duas Novas Espécies do Gênero Mesoblattinopsis (Insecta, Blattoptera, Blattidae) na Formação Santana, Cretáceo do Nordeste do Brasil. *XV Congresso Brasileiro de Paleontologia, 1997, São Pedro, Boletim de Resumos 1*: 58.
- 1998a. An attempt at taphonomic simulation in blattopteras from Araripe (Lower Cretaceous), from the North East of Brazil. *1st Paleoentomological Conference, Abstracts, 30 August–4 September 1998, Moscow*: 25.
- 1998b. An Attempt at taphonomic simulation in Blattopteras from Araripe (Lower Cretaceous) from the NE of Brazil. *AMBA projects International 98*: 99–101.
- 2000. Novos Blattodeas (Insecta, Blattopteroidea) da Formação Santana, Cretáceo Inferior do Nordeste do Brasil. *Revista Universidade de Guarulhos - Ciências Exatas e Tecnológicas 5(6)*: 25–35.
- Menon, F. 2005. New record of Tettigarctidae (Insecta, Hemiptera, Cicadoidea) from the Lower Cretaceous of Brazil. *Zootaxa 1087*: 53–58.
- and Heads, S. W. 2005. New species of Palaeontinidae (Insecta: Cicadomorpha) from the Lower Cretaceous Crato Formation of Brazil. *Stuttgarter Beiträge zur Naturkunde, Serie B 357*: 1–11.
- , Heads, S. W. and Martill, D. M. 2005. New Palaeontinidae (Insecta: Cicadomorpha) from the Lower Cretaceous Crato Formation of Brazil. *Cretaceous Research 26*: 837–844.
- Michener, C. D. 2000. *The Bees of the World*. Baltimore, MD: The John Hopkins University Press.
- Mickoleit, G. 1973. Über den Ovipositor der Neuropteroidea und Coleoptera und seine phylogenetische Bedeutung (Insecta, Holometabola). *Zeitschrift für Morphologie der Tiere 74*: 37–64.
- Misof, B. and Rickert, A. 1999a. A molecular approach to the phylogeny of extant Odonata. *Zoology 102* (suppl. II): 9.
- and Rickert, A. 1999b. A molecular phylogeny of Anisoptera. *Zoology 102* (suppl. II): 9.
- Mostovski, M. B. and Jarzembowski, E. A. 2000. The first brachycerous flies (Diptera: Rhagionidae) from the Lower Jurassic of Gondwana. *Paleontological Journal 34* (suppl. 3): 367–369.
- and Martínez-Delclós, X. 2000. New Nemestrinoidea (Diptera: Brachycera) from the Upper Jurassic – Lower Cretaceous of Eurasia, taxonomy and paleobiology. *Entomological Problems 31*: 137–148.
- , Jarzembowski, E. A. and Coram, R. A. 2003. Horseflies and athericids (Diptera: Tabanidae, Athericidae) from the Lower Cretaceous of England and Transbaikalia. *Paleontological Journal 37*: 162–169.

- Moulds, M. S. 1990. *Australian Cicadas*. Kensington, NSW: New South Wales University Press.
- 2005. An appraisal of the higher classification of cicadas (Hemiptera: Cicadoidea) with special reference to the Australian Fauna. *Records of the Australian Museum* **57**: 375–446.
- Nagatomi, A. and Yang, D. 1998. A review of extinct Mesozoic genera and families of Brachycera (Insecta, Diptera, Orthorrhapha). *Entomologist's Monthly Magazine* **134**: 95–192.
- and Rozkosny, R. 2000. Xylomyidae, pp. 369–378. In Papp, L. and Darvas, B. (eds), *Manual of Palaearctic Diptera*, vol. 2. Budapest: Science Herald.
- Nagy, C. G. 1969. A new taxon of the family Heterogynidae Latreille (Hym. Aculeata). *Entomologische Mitteilungen aus dem Zoologischen Staatsinstitut und Zoologischen Museum Hamburg* **64**: 299–303.
- Neboiss, A. 1960. On the family Cupedidae, Coleoptera. *Proceedings of the Royal Society of Victoria* **72**: 12–20.
- Nel, A. 1996. Un Tettigarctidae fossile du Lias Européen (Cicadomorpha, Cicadoidea, Tettigarctidae). *Ecole Pratique des Hautes Etudes, Biologie et Evolution des Insectes* **9**: 83–94.
- and Paicheler, J.-Cl. 1992. Les Heteroptera aquatiques fossiles, état actuel des connaissances (Heteroptera: Nepomorpha et Gerromorpha). *Entomologica Gallica* **3**: 159–182.
- and Martínez-Delclòs, X. 1993. Essai de révision des Aeschnidioidea (Insecta, Odonata, Anisoptera). *Cahiers de Paléontologie* **1993**: 1–99.
- and Paicheler, J.-C. 1993. Les Isoptera fossiles, pp. 103–179. In Nel, A., Martínez-Delclòs, X. and Paicheler, J.-C. (eds), *Essai de révision des Aeschnidioidea [Insecta, Odonata, Anisoptera] / Les Isoptera fossiles [Insecta, Dictyoptera]*. Paris: CNRS Editions (Cahiers de Paléontologie).
- and Escuillé, F. 1994. A new dragonfly from the Lower Cretaceous of Brazil. *Palaeontology* **37**: 923–930.
- and Paicheler, J.-C. 1994a. Les Gomphidae fossiles. Un inventaire critique (Odonata: Gomphidae). *Annales de la Société Entomologie de France* (n.s.) **30**: 55–77.
- and Paicheler, J.-C. 1994b. Les Libelluloidea fossiles autres que Libellulidae. Un inventaire critique (Odonata, Corduliidae, Macromiidae, Synthemistidae, Chlorogomphidae et Mesophlebiidae). *Nouvelle Revue d'Entomologie* (N.S.) **11**: 321–334.
- and Popov, Yu. A. 2000. The oldest known fossil Hydrometridae from the Lower Cretaceous of Brazil (Heteroptera: Gerromorpha). *Journal of Natural History* **34**: 2315–2322.
- and Waller, A. 2006. A giant water bug from the Lower Cretaceous Crato Formation of Brazil (Heteroptera: Belostomatidae: Lethocerinae). *Zootaxa* **1220**: 63–68.
- Sémeria, Y. and Martins-Neto, R. G. 1990. Un Raphidioptera fossile du Crétacé inférieur du Brésil (Neuropteroidea). *Neuroptera International* **6**: 27–37.
- , Martínez-Delclòs, Paicheler, J.-C. and Henrotay, M. 1993. Les “Anisozygoptera” fossiles. Phylogénie et classification. (Odonata). *Martinia, Numéro hors-série* **3**: 1–311.
- , Bechly, G., Jarzembski, E. A. and Martínez-Delclòs, X. 1998. A revision of the fossil petalurid dragonflies (Insecta: Odonata: Anisoptera: Petalurida). *Paleontologia Lombarda* **10**: 1–68.
- , Zarbout, Barale, G. and Philippe, M. 1998. *Liassotettigarcta africana* sp. n. (Auchenorrhyncha: Cicadoidea: Tettigarctidae), the first Mesozoic insect from Tunisia. *European Journal of Entomology* **95**: 593–598.

- , Azar, D., Martínez-Delclós, X. and Makhoul, E. 2004. A new Upper Cretaceous species of *Chresmoda* from Lebanon – a latest representative of Chresmodidae (Insecta: Polyneoptera inc. sed.): first record of homeotic mutations in the fossil record of insects. *European Journal of Entomology* **101**: 145–151.
- , Perrault, G., Perrichot, V. and Néraudeau, D. 2004. The oldest ants in the Lower Cretaceous amber of Charente – Maritime (SW France) (Insecta: Hymenoptera: Formicidae). *Geologica Acta* **2**: 23–29.
- , Bechly, G., Garouste, R., Pohl, B. and Escuillié, F. 2005. A new extraordinary neuropterid family from the Lower Cretaceous Crato Formation of Brazil: a new insect order? (Insecta, Neuroptera). *Cretaceous Research* **26**: 845–852.
- , Martínez-Delclós, X. and Hutin, A.. 2005. Mesozoic chrysopid-like Planipennia: a phylogenetic approach (Insecta: Neuroptera). *Annales de la Société Entomologie de France (n.s.)* **41**: 29–69.
- Newton, Jr, A. F. and Thayer, M. K. 1995. Protopselaphinae new subfamily for *Protopselaphus* new genus from Malaysia, with a phylogenetic analysis and review of the Omaliine Group of Staphylinidae including Pselaphidae (Coleoptera), pp. 219–320. In Pakaluk, J. and Ślipiński, S. A. (eds), *Biology, Phylogeny, and Classification of Coleoptera*, vol 1. Warsaw: Muzeum I Instytut Zoologii PAN.
- Nikolajev, G. V. 1992. Taxonomic criteria and generic composition of mesozoic lamellicorn beetles (Coleoptera, Scarabaeidae). *Paleontological Journal* **26**: 96–111.
- Novokschonov, V. G. 1997. *Early Evolution of Scorpionflies (Insecta: Panorpida)*. Moscow: Nauka.
- 2002. Order Panorpida Latreille, 1802. The scorpionflies, pp. 194–199. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
- O'Brien, L. B. 2002. The wild wonderful world of Fulgoromorpha, pp. 83–102. In Holzinger, W. (ed), *Zikaden – Leafhoppers, Planthoppers and Cicadas (Insecta: Hemiptera: Auchenorrhyncha)*. Denisia 04, Zugleich Kataloge, Neue Folge Nr. 176. Linz: OÖ Landesmuseums.
- Ogden, T. H. and Whiting, M. F. 2003. The problem with the “Paleoptera Problem:” sense and sensitivity. *Cladistics* **19**: 432–442.
- 2005. Phylogeny of Ephemeroptera (mayflies) based on molecular evidence. *Molecular Phylogenetics and Evolution* **37**: 625–643.
- Ohl, M. 1995. Die phylogenetischen Beziehungen von Grabwespen und Bienen: Stand der Forschung, Probleme, Perspektiven. *Mitteilungen der Deutschen Gesellschaft für allgemeine und angewandte Entomologie* **10**: 629–632.
- 1996. Die phylogenetischen Beziehungen der Sphecinae (Hymenoptera: Apoidea: “Sphecidae”) aufgrund morphologischer Merkmale des Exoskellets. *Zoologische Beiträge (n. f.)* **37**: 3–40.
- 2000. Das phylogenetische System der aculeaten Hymenopteren sensu stricto – Stand der Forschung, Probleme, Perspektiven. *Beiträge der Hymenopterologen-Tagung Stuttgart 2000*: 10–13.
- and Bleidorn, C. 2006. The phylogenetic position of the enigmatic wasp family Heterogynaidae based on molecular data, with description of a new, nocturnal species (Hymenoptera: Apoidea). *Systematic Entomology* **31**: 321–337.
- Olsen, P. E., Remington, C. L., Cornet, B. and Thompson, K. S. 1978. Cyclic change in Late Triassic lacustrine communities. *Science* **201**: 729–733.
- Oman, P. W., Knight, W. J. and Nielson, M. W. 1990. *Leafhoppers (Cicadellidae): a Bibliography, Generic Check-list and Index to the World Literature 1956–1985*. Wallingford: CAB International Institute of Entomology.
- Osten, T. 1987. Ein neuer Fundort von *Proscolia spectator* Day, 1981 (Hymenoptera, Aculeata). *Entomofauna* **8**: 361–365.

- 1988. Die Mundwerkzeuge von *Proscolia spectator* Day (Hymenoptera: Aculeata). Ein Beitrag zur Phylogenie der “Scolioidea”. *Stuttgarter Beiträge zur Naturkunde Ser. A* **414**: 1–30.
- 1993. The enigmatic Genus *Proscolia*, its distribution and phylogenetic relationships. *Biologia Gallo – Hellenica* **20**: 177–182.
- 2001. Beitrag zur Scoliiden – Fauna der Insel Phuket (Thailand) (Hymenoptera, Scoliidae). *Entomofauna* **22**: 433–444.
- 2004. Verbreitungsmuster bei Scoliiden – Ein Beitrag zu ihrer Evolution. *Beiträge der Hymenopterologen-Tagung Stuttgart 2004*: 7–9.
- 2005. Checkliste der Dolchwespen der Welt (Insecta: Hymenoptera, Scoliidae). *Bericht der Naturforschenden Gesellschaft Augsburg* **62**: 1–62.
- Oswald, J. D. 1990. Raphidioptera, pp. 154–163. In Grimaldi, D. A. (ed), *Insects from the Santana Formation, Lower Cretaceous, of Brazil. Bulletin of the American Museum of Natural History* **195**.
- O’Tool, C. and Raw, A. 1991. *Bees of the World*. London: Blanford.
- Ouvrard, D., Campbell, B. C., Bourgois, Th. and Chan, K. L. 2000. 18S rRNA secondary structure and phylogenetic position of Peloridiidae (Insecta, Hemiptera). *Molecular Phylogenetics and Evolution* **16**: 403–417.
- Pate, V. S. L. 1946. The generic names of the Sapygidae and their type species (Hymenoptera: Aculeata). *Entomological News* **57**: 219–221.
- 1947. Neotropical Sapygidae with a conspectus of the family (Hymenoptera: Aculeata). *Acta Zoologica Lilloana* **4**: 393–426.
- Peters, W. L. and Peters, J. G. 2000. Discovery of a new genus of Leptophlebiidae: Leptophlebiinae (Ephemeroptera) in Cretaceous amber from New Jersey, pp. 127–131. In Grimaldi, D. (ed), *Studies on Fossils in Amber, with Particular Reference to the Cretaceous of New Jersey*. Leiden: Backhuys.
- Petrulevičius, J. F. 1998. First hanging fly fossil from South America. *1st Paleoentomological Conference, Abstracts, 30 August–4 September 1998, Moscow*: 34.
- and Martins-Neto, R. G. 2001. A bitacid from the Santana Formation, Lower Cretaceous of Brazil. Legal, ethical and systematic aspects. *Acta Geological Leopoldensia* **24**: 125–127.
- and Martins-Neto, R. G. 2007. A new proterogomphid dragonfly from the Lower Cretaceous of Brazil. *Palaeontology* **50** (in press).
- Pierce, W. D. 1950. Fossil arthropods from onyx marble. *Bulletin of the Southern Californian Academy of Sciences* **49**: 101–104.
- 1951. Fossil arthropods from onyx marble. *Bulletin of the Southern Californian Academy of Sciences* **50**: 44–49.
- Pinto, J. D. 1989. A second new blattoid from the Cretaceous of Brazil. *Resumo das Comunicações do Congresso Brasileiro de Paleontologia* **11**: 295–300.
- and Ornellas, L. P. 1974. New Cretaceous Hemiptera (Insects) from Codo Formation – Northern Brazil. *Anais 28th Congresso Brasileiro do Geologia*: 289–304.
- Pinto, I. R. and Purper, I. 1986. A new blattoid from the Cretaceous of Brazil. *Pesquisas Instituto de Geociências Universidad Federal do Rio Grande do Sul* **18**: 5–10.
- Poinar, J. G. O. and Danforth, B. N. 2006. A fossil bee from Early Cretaceous Burmese amber. *Science* **314**: 614.
- Polegatto, C. M. and Zamboni, J. C. 2001. Inferences regarding the feeding behavior and morphoecological patterns of fossil mayfly nymphs (Insecta Ephemeroptera) from the Lower Cretaceous Santana Formation of northeastern Brazil. *Acta Geologica Leopoldensia* **24**: 145–160.

- Ponomarenko, A. G. 1977. Palaeozoic members of Megaloptera (Insecta). *Paleontologicheskiy Zhurnal* **1977**: 78–86.
- 1986. Scarabaeiformes incertae sedis, pp. 110–112. In Rasnitsyn, A. P. (ed), *Nasekomye v rannemelovyx ekosistemakh Zapadnoy Mongoli. Trudy Sovmestnaya Sovetsko – Mongol'skaya Paleontologicheskaya Ehkspeditsiya* **28**: 1–213.
 - 1987. New Mesozoic water beetles (Insecta, Coleoptera) from Asia. *Paleontological Journal* **21**: 79–92.
 - 1997. New beetles of the Family Cupedidae from the Mesozoic of Mongolia. Ommatini, Mesocupedini, Priacmini. *Paleontological Journal* **31**: 389–399.
 - 2000. New alderflies (Megaloptera: Parasailidae) and glosselytrodeans (Glosselytrodea: Glosselytridae) from the Permian of Mongolia. *Paleontological Journal, suppl. 3* **34**: 309–311.
 - 2003. Ecological evolution of beetles (Insecta: Coleoptera). *Acta Zoologica Cracoviensis* **46** (suppl. Fossil Insects): 319–328.
 - and Martínez-Declòs, X. 2000. New beetles (Insecta: Coleoptera) from the Lower Cretaceous of Spain. *Acta Geologica Hispanica* **35**: 47–52.
- Popham, E. J. 1990. Dermaptera, pp. 69–75. In Grimaldi, D. A. (ed), *Insects from the Santana Formation, Lower Cretaceous, of Brazil. Bulletin of the American Museum of Natural History* **195**.
- Popov, Yu. A. 1971. The historical development of Hemiptera infraorder Nepomorpha (Heteroptera). *Proceedings of the Paleontological Institute, Academy of Sciences, USSR* **129**.
- 1986. Pelorida (=Coleorrhyncha) et Cimicina (=Heteroptera). Insects in Early Cretaceous ecosystems of western Mongolia. *Proceedings of the Joint Soviet-Mongolian Paleontological Expeditions, Nauka, Moscow* **28**: 50–83.
 - 1990. Description of fossil insects. True bugs. Cimicina. In *Late Mesozoic insects of Eastern Transbaikalia. Proceedings of the Paleontological Institute, Academy of Sciences, USSR, Nauka* **239**: 20–39.
 - and Shcherbakov, D. E. 1991. Mesozoic Peloridioidea and their ancestors (Insecta: Hemiptera, Coleorrhyncha). *Geologica et Palaeontologica* **25**: 215–235.
 - and Shcherbakov, D. E. 1996. Origin and evolution of the Coleorrhyncha as shown by the fossil record, pp. 9–30. In Schaefer, C. W. (ed), *Studies on Hemipteran Phylogeny*. Proceedings. Entomological Society of America. Lanham, MD: Thomas Say Publications in Entomology.
 - and Pinto, I. D. 2000. On some mesozoic burrower bugs (Heteroptera: Cydnidae). *Paleontological Journal* **34** (suppl. 3): 298–302.
 - , Dolling, W. R. and Whalley, P. E. S. 1994. British Upper Triassic and Lower Jurassic Heteroptera and Coeleorrhyncha (Insecta: Hemiptera). *Genus* **5**: 307–347.
- Prentice, M. A. 1998. *The Comparative Morphology and Phylogeny of Apoid Wasps (Hymenoptera: Apoidea)*. Dissertation thesis, University of California, Berkley.
- Pritchard, G., McKee, M. H., Pike, E. M., Scrimgeour, G. and Zloty, J. 1993. Did the first insects live in water or in air? *Biological Journal of the Linnean Society* **49**: 31–44.
- Prokop, J., Nel, A. and Hoch, I. 2005. Discovery of the oldest known Pterygota in the Lower Carboniferous of the Upper Silesian Basin in the Czech Republic (Insecta: Archaeorthoptera). *Geobios* **38**, 383–387.
- Pulawski, W. J. and Rasnitsyn, A. P. 2000. *Cretobestiola*, a replacement name for *Bestiola* Pulawski and Rasnitsyn, 1999 (Hymenoptera: Sphecidae). *Acta Geologica Hispanica* **35**: 53.
- Rasnitsyn, A. P. 1977. A new subfamily of scoliid wasps (Hymenoptera, Scoliidae, Proscoliinae). *Zoologicheskii Zhurnal* **56**: 522–529.

- 1986a. Review of the fossil Tippiidae, with description of a new species (Hymenoptera). *Psyche* **93**: 91–101.
- 1988. An outline of evolution of the hymenopterous insects (Order Vespida). *Oriental Insects* **22**: 115–145.
- 1993. Archaeoscoliinae, an extinct subfamily of scoliid wasps (Insecta: Vespa = Hymenoptera: Scoliidae). *Journal of Hymenoptera Research* **2**: 85–96.
- 2002. Infraclass Gryllones, pp. 254–256. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
- and Martínez-Delclòs, X. 1999. New Cretaceous Scoliidae (Vespa = Hymenoptera) from the Lower Cretaceous of Spain and Brazil. *Cretaceous Research* **20**: 767–772.
- and Martínez-Delclòs, X. 2000. Wasps (Insecta: Vespa = Hymenoptera) from the Early Cretaceous of Spain. *Acta Geologica Hispanica* **35**: 65–95.
- and Pritykina, L. N. 2002. Superorder Libellulida Laicharting, 1781. Order Odonata Fabricius, 1792. The dragonflies, pp. 97–106. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
- and Quicke, D. L. J. (eds), 2002. *History of Insects*. Dordrecht: Kluwer.
- and Zherikhin, V. V. 2002. 4. Appendix: alphabetic list of selected insect fossil sites 4.1. impression fossils, pp. 437–444. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
- , Jarzembowski, E. A. and Ross, A. J. 1998. Wasps (Insecta: Vespa = Hymenoptera) from the Purbeck and Wealden (Lower Cretaceous) of Southern England and their biostratigraphical and paleoenvironmental significance. *Cretaceous Research* **19**: 329–391.
- , Pulawski, W. J. and Martínez-Delclòs, X. 1999. Cretaceous digger wasps of the new genus *Bestiola* Pulawski and Rasnitsyn (Hymenoptera: Sphecidae: Angarosphecinae). *Journal of Hymenoptera Research* **8**: 23–34.
- Rehn, A. C. 2003. Phylogenetic analysis of higher-level relationships of Odonata. *Systematic Entomology* **28**: 181–239.
- Ren, D. 1997. First record of fossil stick-insects from China with analyses of some palaeobiological features (Phasmatodea: Hagiphasmatidae fam. nov.). *Acta Zootaxonomica Sinica* **22**: 268–282.
- 1998. Late Jurassic [sic: Early Cretaceous] Brachycera from northeastern China (Insecta: Diptera). *Acta Zootaxonomica Sinica* **23**: 65–83.
- , Lu, L., Guo, Z., Ji, S. and Han, Y. 1995. [*Fauna and Stratigraphy of Jurassic-Cretaceous of Beijing and adjacent areas*]. Beijing: Seismic Publishing House.
- , Yin, J.-C. and Dou, W.-X. 1998a. Late Jurassic palaeontinids (Homoptera: Auchenorrhyncha) from the Hebei and Liaoning provinces in China. *Entomologica Sinica* **5**: 222–232.
- , Yin, J.-C. and Dou, W.-X. 1998b. New planthoppers and froghoppers from the Late Jurassic of Northeast China (Homoptera: Auchenorrhyncha). *Acta Zootaxonomica Sinica* **23**: 281–288.
- Rentz, D. C. F. 1996. *Grasshopper country: the abundant orthopteroid insects of Australia*. Sydney: New South Wales University Press.
- and Su, Y. N. 2003. Orthoptera (grasshoppers, locusts, katydids, crickets), pp. 827–839. In Resh, V. H. and Cardé, R. T. (eds), *Encyclopedia of Insects*. San Diego: Academic Press.
- Ribeiro, G. C. and Martins-Neto, R. G. 1999. A new Tipulidae (Insecta, Diptera) from the Santana Formation (Araripe Basin, Lower Cretaceous, Northeastern Brazil).

- Boletim do 5th Simpósio sobre o Cretáceo do Brasil (UNESP, São Paulo) 1999:* 207–212.
- and Krzemiski, W. 2000. New information on Limoniidae (Diptera: Tipulomorpha) from the Lower Cretaceous Santana Formation (northeastern Brazil). *Polskie Pismo Entomologiczne* **69**: 451–457.
- Richter, S. 2002. The Tetraconata concept: hexapod-crustacean relationships and the phylogeny of Crustacea. *Organisms Diversity & Evolution* **2**: 217–237.
- Riek, E. F. 1974. Upper Triassic insects from the Molteno Formation, South Africa. *Palaeontologica Africana* **17**: 19–31.
- Rohdendorf, B. B. 1977. The rationalization of names of higher taxa in zoology. *Paleontological Journal* **11**: 149–155.
- (ed.) 1991. *Fundamentals of Paleontology 9: Arthropoda, Tracheata, Chelicerata*. New Dehli: Amerind Publishers.
- Rohr, D. M., Boucot, A. J., Miller, J., and Abbott, M. 1986. Oldest termite nest from the Upper Cretaceous of west Texas. *Geology* **14**: 87–88.
- Romero, E. J. 1986. Fossil evidence regarding the evolution of *Nothofagus* Blume. *Annals of the Missouri Botanical Garden* **73**: 276–283.
- Roth, L. M. 1994. The beetle-mimicking cockroach genera *Prosoplecta* and *Areolaria*, with a description of *Tomeisneria furthi* gen. n., sp. n. (Blattellidae: Pseudophyllodromiinae). *Entomologica Scandinavica* **24**: 419–426.
- Rowell, C. H. F. and Flook, P. K. 1998. Phylogeny of the Caelifera and the Orthoptera as derived from ribosomal gene sequences. *Journal of Orthoptera Research* **7**: 147–156.
- Rozkosny, R. 2000. Stratiomyidae, pp. 387–411. In Papp, L. and Darvas, B. (eds), *Manual of Palaearctic Diptera*, vol. 2. Budapest: Science Herald.
- Ruf, M. L., Goodwyn, P. P. and Martins-Neto, R. G. 2005. New Heteroptera (Insecta) from the Santana Formation, Lower Cretaceous (Northeastern Brazil), with description of a new family and new taxa of Naucoridae and Gelastocoridae. *Gaea* **1**: 68–74.
- Rumbucher, K. 1995. Hemerobiidae (Insecta, Planipennia), eine bisher noch nicht entdeckte Familie der Santana Formation aus der brasilianischen Unterkreide. *Bericht der Naturforschenden Gesellschaft Augsburg* **55**: 46–61.
- Sakakibara, A. M. 1979. Sobre algumas espécies brasileiras de *Deois* Fennah, 1948 (Homoptera, Cercopidae). *Revista Brasileira de Biologia* **39**: 9–30.
- Samuelson, G. A. 1994. Pollen consumption and digestion by leaf beetles, pp. 179–183. In Jolivet, P. H., Cox, M. L. and Petitpierre, E. (eds), *Novel Aspects of the Biology of Chrysomelidae*. Dordrecht: Kluwer Academic Publishers.
- Santiago-Blay, J. A. 1994. Paleontology of leaf beetles, pp. 1–68. In Jolivet, H., Cox, M. L. and Petitpierre, E. (eds.), *Novel Aspects of the Biology of Chrysomelidae*. Dordrecht: Kluwer Academic Publishers.
- Santos, M. E. C. 1971. Um novo artrópodo da Formação Areão, Estado de Minas Gerais. *Annales Academiae Brasiliensis Ciencias* **43**: 415–420.
- Schaefer, C. W. (ed.). 1996. *Studies on Hemiptera Phylogeny*. Langham, MD: Thomas Say Publications in Entomology; Proceedings Entomological Society of America.
- Schlee, D. 1969. Morphologie und Symbiose; ihre Beweiskraft für die Verwandtschaftsbeziehungen der Coleorrhyncha (Insecta, Hemiptera). *Stuttgarter Beiträge zur Naturkunde, Serie C* **210**: 1–27.
- Schlüter, T. 1990. Fossil insect localities in Gondwanaland. *Entomologia Generalis* **15**: 61–76.
- Scholtz, C. H. and Chown, S. L. 1995. The evolution of habitat use and diet in the Scarabaeoidea: a phylogenetic approach, pp. 355–374. In Pakaluk, J. and Ślipiński,

- S. A. (eds), *Biology, Phylogeny, and Classification of Coleoptera*, vol. 1. Warsaw: Muzeum I Instytut Zoologii PAN.
- Schuh, R. T. and Slater, J. A. 1995. *True Bugs of the World (Hemiptera: Heteroptera) – Classification and Natural History*. Ithaca: Cornell University Press.
- Scudder, S. H. 1880. *The Devonian Insects of New Brunswick*. Anniversary Memoires of the Boston Society of Natural History.
- Sellick, J. T. C. 1994. Phasmida (stick insect) eggs from the Eocene of Oregon. *Palaeontology* **37**: 913–921.
- 1997. Descriptive terminology of the phasmid egg capsule, with an extended key to the phasmid genera based on egg structure. *Systematic Entomology* **22**: 97–122.
 - 1998. The micropylar plate of the eggs of Phasmida, with a survey of the range of plate form within the order. *Systematic Entomology* **23**: 203–228.
- Sharov, A. G. 1968. Phylogeny of the Orthopteroidea. *Trudy Palaeontologicheskogo Instituta, Nauk* **118**: 1–251.
- Shear, W. A., Bonamo, P. M., Grierson, J. D., et al. 1984. Early land animals in North America. *Science* **224**: 492–494.
- Shcherbakov, D. E. 1986 (1985). Nasekomye v rannemelovykh ekosistemach zapadnoi Mongoli. *Transactions of the joint Soviet–Mongolian Paleontological Expedition* **28**: 47–50.
- 1988. New cicadas (Cicadina) from the later Mesozoic of Transbaikalia. *Paleontological Journal* **4**: 52–63.
 - 1992. The earliest leafhoppers (Hemiptera: Karajassidae n. fam.) from the Jurassic of Karatau. *Neues Jahrbuche für Geologie und Paläontologie, Monatshefte* **1**: 39–51.
 - 1996. Origin and evolution of the Auchenorrhyncha as shown by the fossil record, pp. 31–45. In Schaefer, C. W. (ed.), *Studies on Hemipteran Phylogeny*. Lanham, MD: Thomas Say Entomological Publications. Entomological Society of America.
 - 2000. The most primitive white-flies (Hemiptera, Aleyrodidae, Bernaeinae subfam. nov.) from the Mesozoic of Asia and Burmese amber with an overview of Burmese amber hemipterans. *Bulletin of the Natural History Museum, Geology* **56**: 29–37.
 - 2002. Order Forficulida. The earwigs and protelytropterans, pp. 288–291, 298–301. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
 - 2004. Fossil adult and nymphal Tettigarctidae, pp. 71–73. In Kerzhner, I. M. (ed), *Third European Hemiptera Congress. St. Petersburg, June 8–11, 2004. Abstracts*. St. Petersburg: Russian Academy of Sciences, Department of Biological Sciences, St. Petersburg Scientific Centre, Zoological Institute.
 - and Popov, Yu. A. 2002. 2.2.1.2.5. Superorder Cimicidea Laicharting, 1781 Order Hemiptera Linné, 1758. The Bugs, Cicadas, Plantlice, Scale Insects, etc. (= Cimicida Laicharting, 1781, = Homoptera Leach, 1815 + Heteroptera Latreille, 1810), pp. 143–157. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
- Silvestri, F. 1912. Die Thysanuren des Baltischen Bernsteins. *Schriftreihe der physikalisch-ökonomischen Gesellschaft Königsberg* **53**: 42–66.
- Sinitshenkova, N. D. 1975. Mayfly nymphs of the family Hexagenitidae (Insecta, Ephemeroptera). *Paleontologicheskii Zhurnal* **1975**: 82–86.
- 1986. Mayflies. Ephemera (=Ephemeroptera), In Rasnitsyn, A. P. (ed.), *Insects in the Early Cretaceous Ecosystems of the Western Mongolia. Transactions of the Joint Soviet–Mongolian Paleontological Expedition. Moscow. Nauka* **28**: 45–46.
 - 2000. The first fossil prosopistomatid mayfly from Burmese amber (Ephemeroptera: Prosopistomatidae). *Bulletin of the Natural History Museum of London (Geology)* **56**: 23–26.

- and Coram, R. A. 2002. The first mayfly from the Lower Cretaceous of southern England (Insecta: Ephemeroidea =Ephemeroptera). *Cretaceous Research* **23**: 461–463.
- Marchal-Papier, F., Grauvogel-Stamm, L. and Gall, J.-C. 2005. The Ephemeroidea (Insecta) from the Grès à Voltzia (early Middle Triassic) of the Vosges (NE France). *Paläontologische Zeitschrift* **79**: 377–397.
- Smith, P. E., Evensen, N. M., York, D. et al. 1995. Dates and rates in ancient lakes: ^{40}Ar - ^{39}Ar evidence for an Early Cretaceous age for the Jehol Group, Northeast China. *Canadian Journal of Earth Sciences* **32**: 1426–1431.
- Snodgrass, R. E. 1935. *Principles of Insect Morphology*. New York: McGraw-Hill.
- Sorensen, J. T., Campbell, B. C., Gill, R. J. and Steffen-Campbell, J. D. 1995. Non-monophyly of Auchenorrhyncha ("Homoptera"), based upon 18S rDNA phylogeny: eco-evolutionary and cladistic implications within pre-Heteropterodea Hemiptera (s.l.) and a proposal for new monophyletic sub-orders. *Pan-Pacific Entomologist* **71**: 31–60.
- Spahr, U. 1987. Ergänzungen und Berichtigungen zu R. Keilbachs Bibliographie und Liste der Bernsteinfossiliien-Ordnung Hymenoptera. *Stuttgarter Beiträge zur Naturkunde, Serie B* **127**: 1–121.
- Staniczek, A. H. 1997. The morphology of *Siphlaenigma janae* Penniket (Ephemeroptera, Siphlaenigmatidae), and its significance for the ground plan of the Baetoidea, pp. 536–549. In Landolt, P. and Sartori, M. (eds), *Ephemeroptera and Plecoptera: Biology-Ecology-Systematics*. Freibourg: Mauron + Tinguely & Lachat SA.
- 2000. The mandible of Silverfish (Insecta: Zygentoma) and Mayflies (Ephemeroptera): its morphology and phylogenetic significance. *Zoologischer Anzeiger* **239**: 147–178.
- 2003. New fossil mayflies from Dominican amber (Insecta: Ephemeroptera: Leptophlebiidae: Atalophlebiinae). *Stuttgarter Beiträge zur Naturkunde (B)* **341**: 1–22.
- and Bechly, G. 2002. First fossil record of the mayfly family Baetiscidae from Baltic amber (Insecta: Ephemeroptera). *Stuttgarter Beiträge zur Naturkunde (B)* **322**: 1–11.
- Stuckenberg, B. R. 2001. Pruning the tree: a critical review of classifications of the Homeodactyla (Diptera, Brachycera), with new perspectives and an alternative classification. *Studia Dipterologica* **8**: 3–41.
- Sturm, H. 1998. Erstnachweis fischchenartiger Insekten (Zygentoma, Insecta) für das Mesozoikum (Untere Kreide, Brasilien). *Senckenbergiana Lethaea* **78**: 135–140.
- and Mendes, L. F. 1998. Two new species of Nicoletiidae (Zygentoma, 'Apterygota', Insecta) in Dominican amber. *American Museum Novitates* **3226**: 1–11.
- and Machida, R. 2001. *Archaeognatha*. Handbuch der Biologie Band 37/ IV. Berlin: Akademische Verlagsgesellschaft Athenaion.
- Svensson, G. J. and Whiting, M. F. 2004. Phylogeny of Mantodea based on molecular data: evolution of a charismatic predator. *Systematic Entomology* **29**: 359–370.
- Sweet, M. H. 1996. Comparative external morphology of the pregenital abdomen of the Hemiptera, pp. 119–158. In Schaefer, C. W. (ed), *Studies on Hemiptera Phylogeny*. Langham, MD: Thomas Say Publications in Entomology; Proceedings Entomological Society of America.
- Swisher III, C. C., Wang, Y.-Q., Wang, X.-L., Wang, Xu, X. and Wang, Y. 1999. Cretaceous age for the feathered dinosaurs from Liaoning, China. *Nature* **400**: 58–61.
- Szwedo, J. 2002. Amber and amber inclusions of planthoppers, leafhoppers and their relatives (Hemiptera, Archaeorrhyncha et Clypaeorrhyncha), pp. 37–56. In Holzinger, W. (ed.), *Zikaden – Leafhoppers, Planthoppers and Cicadas (Insecta:*

- Hemiptera: Auchenorrhyncha). Denisia*, 04, Zugleich Kataloge, Neue Folge Nr. 176, Linz: OÖ. Landesmuseums.
- 2004a. A new genus and six new species of ground-dwelling leafhoppers from Chile and New Zealand (Hemiptera: Cicadomorpha: Myerslopiidae). *Zootaxa* **424**: 1–20.
 - 2004b. An annotated checklist of Myerslopiidae with notes on the distribution and origin of the group (Hemiptera: Cicadomorpha). *Zootaxa* **425**: 1–15.
 - 2004c. *Niryasaburnia* gen. nov. for ‘*Liburnia*’ burmitina Cockerell, 1917 from Burmese amber (Hemiptera, Fulgoromorpha: Achilidae). *Journal of Systematic Palaeontology* **2**: 105–107.
 - 2006. A new genus *Waghilde* gen. nov. representing a new tribe of the planthopper family Achilidae from the Eocene Baltic amber (Hemiptera: Fulgoromorpha). *Annales Zoologici* **56**: 167–174.
 - and Webb, M. D. 1999. First fossil representatives of Aetalionidae (Homoptera, Membracoidea) from Oligocene / Miocene ambers of the New World. *10th International Auchenorrhyncha Congress*, Abstracts of Talks and Posters, 6–10 September 1999, Cardiff, Wales: 3–4.
 - and Stroiński A. 2001. *Ptychogroehnia reducta* gen. and sp. n. of the fossil tribe Ptychoptilini from the Eocene Baltic amber (Hemiptera: Fulgoroidea: Achilidae). *Annales Zoologici* **51**: 95–101.
 - , Bourgoin, Th. and Lefebvre, F. 2004. *Fossil Planthoppers (Hemiptera: Fulgoromorpha) of the World. An Annotated Catalogue with Notes on Hemiptera Classification*. Warsaw: Studio 1.
 - Tan, J.-J., Ren, D. and Liu, M. 2005. New ommatids from the Late Jurassic of western Liaoning, China (Coleoptera: Archostemata). *Insect Science* **12**: 207–216.
 - Tanai, T. 1986. Phytogeographic and phylogenetic history of the genus *Nothofagus* Bl. (Fagaceae) in the Southern Hemisphere. *Journal of the Faculty of Science Hokkaido University* **21**: 505–582.
 - Thorne, B. L., Grimaldi, D. A. and Krishna, K. 2000. Early fossil history of the termites, pp. 77–94. In Abe, T., Bignell, D. E. and Higashi, M. (eds), *Termites – Evolution, Sociality, Symbiosis, Ecology*. Dordrecht: Kluwer.
 - and Traniello, J. F. A. 2002. Comparative social biology of basal taxa of ants and termites. *Annual Review of Entomology* **48**: 283–306.
 - Tilgner, E. H. 2000. The fossil record of Phasmida (Insecta: Neoptera). *Insect Systematics and Evolution* **31**: 473–480.
 - Tillyard, R. J. 1922. Mesozoic insects of Queensland. Nr. 9. Orthoptera and additioins to Protorrhoptera, Odonata, Hemiptera and Plannipennia. *Proceedings of the Linnaean Society of New South Wales* **47**: 447–470.
 - 1932. Kansas Permian Insects. Part 15. The Order Plectoptera. *American Journal of Science* **223**: 97–134, 237–272.
 - Tobias, V. I. 1965. Contribution to the knowledge of the family Fedtschenkiidae (Hymenoptera, Sapygoidea). *Zoologicheskii Zhurnal* **XLIV**: 706–715.
 - Trueman, J. W. H. 1996. A preliminary cladistic analysis of odonate wing venation. *Odonatologica* **25**: 59–72.
 - Tshernova, O. A. 1962. Otrjad Ephemeroptera. Podenki, pp. 55–64. In Rodendorf, B. B. (ed), *Osnovy paleontologii*. Moscow: Akademija Nauk SSSR.
 - 1965. Some fossil mayflies (Ephemeroptera Misthodotidae) from Permian-beds of the Urals. *Entomologicheskoe Obozrenie* **44**: 353–361.
 - 1967. Mayfly of the Recent family in Jurassic deposits of Transbaikalia (Ephemeroptera, Siphlonuridae). *Entomologicheskoe Obozrenie* **46**: 322–326.

- 1970. On the classification of Fossil and Recent Ephemeroptera. *Entomological Review* **49**: 71–81.
- and Sinitshenkova, N. D. 1974. A new fossil genus and species of the mayfly family Hexagenitidae (Ephemeroptera) from the South of the European part of the USSR and its connection with Recent mayflies. *Entomologicheskoe Obozrenie* **53**: 130–136.
- Ueda, K. 1997. A new palaeontinid species from the Lower Cretaceous of Brazil (Homoptera: Palaeontinidae). *Bulletin of the Kitakyushu Museum of Natural History* **16**: 99–104.
- Von Brischke, D. 1886. Die Hymenopteren des Bernsteins. *Schriften der Naturforschenden Gesellschaft in Danzig (N.F.)* **6**: 278–279.
- Vršanský, P. 1999a. Lower Cretaceous Blattaria, pp. 167–176. In Vršanský, P. (ed), *Proceedings of the First Palaeoentomological Conference Moscow 1998*. Bratislava: AMBA Projects International.
- 1999b. Two new species of Blattaria (Insecta) from the Lower Cretaceous of Asia, with comments on the origin and phylogenetic position of the families Polyphagidae and Blattulidae. *Entomological Problems* **30**: 85–91.
- 2002. Origin and evolution of mantises. *AMBA Projekty* **6(1)**: 1–16.
- 2003a. Umenocoleoidea – an amazing lineage of aberrant insects (Insecta, Blattaria). *AMBA Projekty* **7**: 1–32.
- 2003b. Unique assemblage of Dictyoptera (Insecta – Blattaria, Mantodea, Isoptera) from the Lower Cretaceous of Bon Tsagaan Nuur in Mongolia. *Entomological Problems* **33(1–2)**, 119–151.
- 2004. Cretaceous Gondwanian Cockroaches (Insecta: Blattaria). *Entomological Problems* **34(1–2)**: 49–54.
- 2005. Evolúcia druhohorných švábov. Ph.D.-thesis. Geological Institute, Slovak Academy of Sciences. 40 pp + 14 supplements.
- , Vishniakova, V. N. and Rasnitsyn, A. P. 2002a. Order Blattida Latreille, 1810. The cockroaches, pp. 263–269. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
- , Mostovski, M. B., Bazylev, B. A. and Bugdaeva, E. 2002b. Early Cretaceous climate changes suggested on the basis of cockroach wing variations. *Proceedings of the XVII Congress of Carpathian–Balkan Geological Association*, 1–4 September 2002 (CD), Bratislava. *Geologica Carpathica* **33** (special issue).
- Vulcano, M. A. 1985. *Cretaceosimulium araripensis* gen. et sp. nov. da Chapada do Araripe, Ceará, Brasil (Diptera: Simuliidae). *Resumos do Congresso Brasileiro de Zoologia* **12**: 107.
- and Pereira, F. S. 1975. Cupesidae (Coleoptera). *Arquivos do Instituto Biológico São Paulo* **42**: 31–68.
- and Pereira, F. S. 1987. Entomofauna fóssil da Chapada do Araripe, Ceará, Brasil – *Cretaceimelittomoides cearensis* gen. nov., sp. nov. (Coleoptera: Pyrochroidae). *X Congreso Brasileiro de Paleontología, 1987, Rio de Janeiro. Resumo das Comunicações*: 27.
- Waller, A., Caussanel, C. and Jamet, C. 1996. Variation morphologique des pieces buccales chez quelques Dermapteres. *Bulletin de la Societe entomologique de France* **101**: 523–533.
- Wang, B., Zhang, H.-C. and Fang, Y. 2006a. Some Jurassic Palaeontinidae (Insecta, Hemiptera) from Daohugou, Inner Mongolia, China. *Palaeoworld* **15**: 115–125.
- , Zhang, H. C. and Fang, Y. 2006b. *Gansucossus*, a replacement name for *Yumenia* Hong, 1982 (Insecta, Hemiptera, Palaeontinidae), with description of a new genus. *Zootaxa* **1268**: 59–68.

- , Zhang, H. C., Fang, Y. and Zhang, Z. 2006c. A new genus and species of Palaeontinidae (Insecta: Hemiptera) from the Middle Jurassic of Daohugou, China. *Annales Zoologici* **56**: 757–762.
- , Zhang, H. C. and Fang, Y. 2007. Middle Jurassic Palaeontinidae (Insecta, Hemiptera) from Daohugou of China. *Alavesia* **1**: 89–104.
- Wang, W., Zhang, L., Zheng, S. et al. 2004. A new study on the stratotype and biostratigraphy of the Yixian stage in Yixian-Beipiao region, Liaoning – establishment and study of stratotypes of the Yixian stage. *Acta Geologica Sinica* **78**: 433–447.
- , Zhang, L., Zheng, S. et al. 2005. The age of the Yixian stage and the boundary of Jurassic-Cretaceous – the establishment and study of stratotypes of the Yixian stages. *Geological Review* **51**: 234–242.
- Wang, Y. and Ren, D. 2006. Middle Jurassic *Pseudococcus* fossils from Daohugou, Inner Mongolia in China (Homoptera, Palaeontinidae). *Acta Zootaxonomica Sinica* **31**: 289–293.
- , Ren, D. and Shih, C.-K. 2007a. Discovery of Middle Jurassic Palaeontinids from Inner Mongolia, China (Homoptera: Palaeontinidae). *Progress in Natural Science* **17**: 112–116.
- , Ren, D. and Shih, C.-K. 2007b. New discovery of palaeontinid fossils from the Middle Jurassic in Daohugou, Inner Mongolia (Homoptera, Palaeontinidae). *Science in China, Series D: Earth Sciences* **50**: 481–486.
- Wappler, T., Hinsken, S., Brocks, J. J., Wetzel, A. and Meyer, C. A. 2005. A fossil sawfly of the genus *Athalia* (Hymenoptera: Tenthredinidae) from the Eocene-Oligocene boundary of Altkirch, France. *Palevol* **4**: 7–16.
- Weber, H. 1933. *Lehrbuch der Entomologie*. Jena: Gustav Fischer.
- Weitschat, W. and Wichard, W. 1998. *Atlas der Pflanzen und Tiere im Baltischen Bernstein*. Munich: Pfeil.
- Westfall, M. J. 1980. Cretaceous dragonfly discovery in Brazil. *Selsysia* **9**: 22.
- Weyenbergh, H. 1874. Varia zoologica et paleontologica. *Periodico Zoologico, Sociedad Entomologica Argentina* **1**: 77–111.
- Whalley, P. E. S. 1983. A survey of recent and fossil cicadas (Insecta, Hemiptera–Homoptera) in Britain. *Bulletin of the British Museum of Natural History (Geology)* **37**: 139–47.
- and Jarzembowksi, E. A. 1985. Fossil insects from the Lithographic Limestone of Montsech (Late Jurassic–Early Cretaceous), Lerida Province, Spain. *Bulletin of the British Museum of Natural History (Geology)* **38**: 381–412.
- Wheeler, W. C., Whiting, M., Wheeler, Q. D., and Carpenter, J. M. 2001. The phylogeny of the extant hexapod orders. *Cladistics* **17**: 113–169, 403–404.
- Wickham, H. F. 1914. Twenty new Coleoptera from the Florissant shales. *Transactions of the Entomological Society of America* **40**: 257–270.
- Wieland, F. 2006. The cervical sclerites of Mantodea discussed in the context of dictyopteran phylogeny (Insecta: Dictyoptera). *Entomologische Abhandlungen* **63**: 51–76.
- Wighton, D. C. 1987. *Gomphaeschna obliqua* spec. nov., a new species of Gomphaeschninae from the Lower Cretaceous of northeastern Brazil (Anisoptera, Aeshnidae). *Odonatologica* **16**: 311–314.
- 1988. Odonate fossils from the Lower Cretaceous of Northeastern Brazil. *Proceeding of the 18th International Congress on Entomology*, Vancouver, Abstracts: 64.
- Wilf, P., Labandeira, C. C., Kress, W. J., et al. 2000. Timing the radiations of leaf beetles: hispines on gingers from Latest Cretaceous to Recent. *Science* **289**: 291–294.

- Willmann, R. 1978. *Mecoptera* (Insecta, Holometabola). In Westphal, F. (ed), *Fossilium Catalogus Animalia, Pars 124*. The Hague: Junk.
- 1987. The phylogenetic system of the Mecoptera. *Systematic Entomology* **12**: 519–524.
- 1989. Evolution und Phylogenetisches System der Mecoptera (Insecta: Holometabola). *Abhandlungen Senckenberg Naturforschungsinstitut, Gesellschaft* **544**: 1–153.
- 1994. Raphidiodea aus dem Lias und die Phylogenie der Kamelhalsfliegen (Insecta: Holometabola). *Paläontologische Zeitschrift* **68**: 167–197.
- 1999. The Upper Carboniferous *Lithoneura lameerei* (Insecta, Ephemeroptera?). *Paläontologische Zeitschrift* **73**: 289–302.
- 2003. Die phylogenetischen Beziehungen der Insecta: offene Fragen und Probleme. *Verhandlungen Westdeutscher Entomologentag* **2003**: 1–64.
- 2006. Die Stammgruppenvertreter der Ephemeroptera und ihre Systematik (Insecta: Ephemeroidea, = Ephemeroptera, = Plectoptera; Eintagsfliegen). *Species, Phylogeny & Evolution* **1**: 23–31.
- Wilson, E. O. 1971. *The Insect Societies*, pp. 31–37. Cambridge, MA: Harvard University Press.
- 1985. Ants from the Cretaceous and Eocene Amber of North America. *Psyche* **92**: 205–216.
- , Carpenter, F. M. and Brown, W. L. 1967. The first Mesozoic Ants, with the description of a new Subfamily. *Psyche* **74**: 1–19.
- Wilson, H. M., Dunlop, J. A., Martill, D. M. and Selden, P. A. 2000. Filling the Mesozoic gap: Terrestrial arthropods from the Crato Formation of Brazil. *Palaeontological Association 44th Annual Meeting, Abstracts and Programme*, University of Edinburgh: **38**.
- and Martill, D. M. 2001. A new Japygid Dipluran from the Lower Cretaceous of Brazil. *Palaeontology* **44**: 1025–1031.
- Woodley, N. E. 1986. Parhadrestiinae, a new subfamily for *Parhadrestia* James and *Cretaceogaster* Teskey (Diptera: Stratiomyidae). *Systematic Entomology* **11**: 377–387.
- Wygodzinsky, P. 1961. On a surviving representative of the Lepidotrichidae (Thysanura). *Annals of the Entomological Society of America* **54**: 621–627.
- Yang, C.-T. 2002. Preliminary thoughts on the phylogeny of Coleorrhyncha-Heteroptera (Hemiptera). *Formosan Entomologist* **22**: 297–305.
- Yang, C.-T. and Chang, T.-Y. 2000. *The External Genitalia of Hemiptera (Homoptera – Heteroptera)*. Taichung, Taiwan: Shih Way Publishers.
- Yeates, D. K. 1999. Congruence and controversy: toward a higher-level phylogeny of diptera. *Annual Review of Entomology* **44**: 397–428.
- and Irwin, M. E. 1996. Apioceridae (Insecta: Diptera): cladistic reappraisal and biogeography. *Zoological Journal of the Linnean Society* **116**: 247–301.
- Yeh, W.-B., Yang, C.-T. and Hui, C.-F. 2005. A molecular phylogeny of planthoppers (Hemiptera: Fulgoroidea) inferred from mitochondrial 16S rDNA sequences. *Zoological Studies* **44**: 519–535.
- Yildirim, E. and Gusenleitner, J. 2001. Sapygidae (Hymenoptera, Aculeata) of Turkey. *Linzer biologische Beiträge* **33**: 671–675.
- Zamboni, J. C. 2001. Contribution to the knowledge of the aquatic paleoentomofauna from Santana Formation (Araripe Basin, Lower Cretaceous, Northeast Brazil) with description of new taxa. *Acta Geologica Leopoldensia* **24**: 129–135.
- Zeuner, F. E. 1944. Notes on Eocene Homoptera from the Isle of Mull, Scotland. *Annals and Magazine of Natural History* **11**: 110–117.

- Zhang, H. C. 1997. Jurassic palaeotinids from Karamai, Xinjiang, with a discussion of Palaeotinidae (Homoptera: Palaeotinidae) in China. *Entomologia Sinica* **4**: 312–323.
- , Rasnitsyn, A. P. and Zhang, J. 2002. The oldest known scoliid wasps (Insecta, Hymenoptera, Scoliidae) from Jehol biota of western Liaoning, China. *Cretaceous Research* **23**: 77–86.
- Zhang, J.-F. 1994. Discovery of primitive earwigs (Insecta) from Late Jurassic of Laiyang, Shandong and its significance. *Acta Palaeontologica Sinica* **33**: 229–245.
- Zhang, Z.-J. 2002. New Early Cretaceous lalacid from Jingxi Basin of Beijing, China. *Acta Zootaxonomica Sinica* **27**: 20–23.
- Zherikhin, V. V. and Gratshev, V. G. 2004. Fossil Curculionoid Beetles (Coleoptera, Curculionoidea) from the Lower Cretaceous of Northeastern Brazil. *Paleontological Journal* **38**: 528–537.
- Zhou, C.-F. and Peters, J. G. 2003. The nymph of *Siphluriscus chinensis* and additional imaginal description: a living mayfly with Jurassic origins (Siphluriscidae new family: Ephemeroptera). *Florida Entomologist* **86**: 345–352.